

EXPRESS MAIL NO. EL491587139US
ATTORNEY DOCKET NO. 21099.0074U2
PATENT

TO ALL WHOM IT MAY CONCERN:

Be it known that we, RICHARD B. MEAGHER, a citizen of the United States of America, residing at 635 Sandstone Drive, Athens, GA 30605, U.S.A., and VINCE LATERZA, a citizen of the United States of America, residing at 610 Webster Drive, Decatur, GA 30033, have invented new and useful improvements in

RAPID PRODUCTION OF MONOCLONAL ANTIBODIES

for which the following is a specification.

RAPID PRODUCTION OF MONOCLONAL ANTIBODIES

This application claims priority to U.S. Provisional Application No. 60/270,322 filed on February 20, 2001, which is hereby incorporated by this reference in its
5 entirety.

FIELD OF THE INVENTION

The present invention relates to genetically altered hybridomas, myelomas and B cells. The invention also relates to utilizing genetically altered hybridomas,
10 myelomas and B cells in methods of making monoclonal antibodies.

BACKGROUND

Major efforts in functional genomics and proteomics are creating an
15 unprecedented demand for monoclonal antibodies to be used for protein function studies. Monoclonal antibodies can be used in every stage of research involving protein target discovery and characterization including purification, quantification, and organ and cellular localization. Recent advances in proteomics are creating a need for large numbers of antibodies for use in high throughput studies and on protein chips.
20 Monoclonal antibodies have been used for decades as key reagents in clinical diagnostics and they are emerging as an important new class of therapeutics agents.

Hybridoma technology is the most commonly used method for accessing monoclonal antibodies. Monoclonal antibodies are secreted from hybridoma cells, created by fusing normal antibody producing splenic B-cells with immortal myeloma
25 cells or other immortal cells. Hybridoma production has changed little since its inception 26 years ago (Kohler and Milstein, 1975).

A typical protocol for hybridoma generation involves: (i) immunizing an animal (e.g., mouse, rat or rabbit) with a purified protein antigen; (ii) harvesting antibody producing B-cells, typically from the spleen; (iii) fusing B-cells with a non-secretory

myeloma cell line deficient for the enzyme hypoxanthine guanine phosphoribosyl transferase (e.g., x63-Ag 8.653 from a BALB/c mouse strain); (iv) growing hybridoma cells in a selection medium containing hypoxanthine, aminopterin and thymidine (HAT) and (v) screening for cells that produce the desired antibody and (vi) limit
5 dilution cloning to obtain a homogenous cell line that secretes the antibody (Antczak, 1982).

Conventional hybridoma technology does not allow researchers to access large numbers of antibodies to different antigens or large numbers of antibodies to a single target antigen in an efficient manner. Hybridoma cell cloning by limit dilution is
10 perhaps the most problematic, time consuming and labor intensive step in generating monoclonal antibodies (O'Reilly et.al.,1998). In this step, cells are repeatedly diluted out to low cell numbers and their supernatants assayed for secreted monoclonal antibody. Screening must be artfully performed to ensure that desired hybridomas are not lost. In the case of rapidly growing hybridomas, cells may die in the microtiter
15 wells from exhaustion of nutrients if they are not moved to larger vessels or fresh medium quickly. Also, in typical wells containing several hybridomas, undesirable hybridomas may continuously overgrow desired hybridomas. This can cause the limit dilution step to be extended weeks or months and may even result in loss of important hybridomas. If the hybridomas have not grown to a reasonable size by the time of
20 assay, they may not have produced sufficient antibody for detection. Therefore, a time for screening supernatants must be chosen carefully. The available "window" for initial screening is not large and usually extends over two to three days (Antczak, 1982). Once started, the limit dilution isolation of pure cell lines typically goes on for 3-4 weeks for any one hybridoma.

25 There is a need for more rapid methods of isolating desired hybridoma cells. At least two laboratories attempted to sort normal hybridomas based on traces of surface presentation of antibody (Parks et al., 1979; Meilhoc et al., 1989). They showed that a subset of hybridoma cells in any population presented a small but measurable number (~ 20) of surface antibody molecules. This was enough to sort these cells when labeled

with antigen, if the antigen was coupled to highly fluorescent microspheres to increase the fluorescence signal. Even with these highly fluorescent spheres the signal was only a few-fold above the background.

There is a need for a significant increase in the presentation of surface antibody on

5 hybridoma cells as well as a need for a significant increase in the percentage of hybridoma cells presenting surface antibody in any population to enable rapid screening. The present invention provides both by providing DISH (Direct Screening of Hybridoma Cells). The DISH technology of the present invention provides a simple, rapid and reliable selection of hybridoma cells that may be accomplished in a matter of

10 hours instead of weeks. DISH provides several significant improvements over conventional hybridoma technology that allow researchers to access much larger repertoires of antibodies in an efficient manner. Using current hybridoma protocols approximately 40,000 fusions are typically prepared. A main reason more fusions are not made is the difficulty encountered in isolating desired clones using limit dilution.

15 DISH enables very rapid, high throughput cell selection using fluorescence activated cell sorting (FACS) and other modalities. Since FACS technology permits millions of cells to be sorted in a matter of hours, the number of hybridoma fusions one can screen using DISH technology is orders of magnitude larger than by limit dilution. Significantly, FACS sorting allows for single cell deposition of desired hybridomas into

20 discrete wells. Hence, the problem of desirable, but slow growing cells being lost is eliminated using DISH. Thus, DISH replaces current antibody screening and limit dilution procedures with a rapid, high throughput, selection process. The present invention can also be utilized to provide populations of plasma cells that surface present adequate immunoglobulin to enable high throughput fluorescence activated cell sorting

25 technology to be used to determine whether single plasma cells produce immunoglobulin that reacts with target antigens.

SUMMARY OF THE INVENTION

5 The present invention provides a population of hybridoma cells wherein greater than 15 % of the cells in the population express monoclonal antibody that is bound to the cell surface.

Also provided by the present invention is a population of hybridoma cells wherein greater than 25 % of the cells in the population express monoclonal antibody that is bound to the cell surface.

10 Further provided is a population of hybridoma cells wherein greater than 50% of the cells in the population express monoclonal antibody that is bound to the cell surface.

Also provided by the present invention is a population of hybridoma cells wherein greater than 75% of the cells in the population express monoclonal antibody that is bound to the cell surface.

15 The present invention also provides a hybridoma cell, wherein greater than twenty monoclonal antibody molecules are expressed and bound to the cell surface.

The present invention further provides a hybridoma cell, wherein greater than fifty monoclonal antibody molecules are expressed and bound to the cell surface.

20 Also provided by the present invention is a hybridoma cell, wherein greater than one hundred monoclonal antibody molecules are expressed and bound to the cell surface.

Further provided by the present invention is a hybridoma cell, wherein greater than two hundred and fifty monoclonal antibody molecules are expressed and bound to the cell surface.

25 Also provided by the present invention is a hybridoma cell, wherein greater than five hundred monoclonal antibody molecules are expressed and bound to the cell surface.

Further provided by the present invention is a population of hybridoma cells wherein greater than 15% of the cells in the population express monoclonal antibody

that is bound to the cell surface and wherein greater than twenty monoclonal antibody molecules are expressed and bound to the cell surface of the cells in the population that express monoclonal antibody.

Also provided by the present invention is a population of hybridoma cells wherein
5 greater than 15% of the cells in the population express monoclonal antibody that is bound to the cell surface and wherein greater than fifty monoclonal antibody molecules are expressed and bound to the cell surface of the cells in the population that express monoclonal antibody.

Further provided by the present invention is a population of hybridoma cells
10 wherein greater than 25% of the cells in the population express monoclonal antibody that is bound to the cell surface and wherein greater than twenty monoclonal antibody molecules are expressed and bound to the cell surface of the cells in the population that express monoclonal antibody.

The present invention also provides a population of hybridoma cells wherein
15 greater than 25% of the cells in the population express monoclonal antibody that is bound to the cell surface and wherein greater than fifty monoclonal antibody molecules are expressed and bound to the cell surface of the cells in the population that express monoclonal antibody.

Also provided by the present invention is a hybridoma cell, wherein from about
20 0.01% to about 10% of the total amount of monoclonal antibody produced by the hybridoma cell is expressed and bound to the cell surface.

Also provided by the present invention is a population of hybridoma cells wherein
greater than 15% of the hybridoma cells in the population express from about 0.01% to about 10% of the total amount of monoclonal antibody produced by the hybridoma
25 cells on the cell surface.

Further provided by the present invention is a population of hybridoma cells wherein greater than 15% of the hybridoma cells in the population express from about 0.01% to about 10% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface, and wherein greater than twenty monoclonal

antibodies are expressed and bound to the cell surface.

Also provided by the present invention is a hybridoma cell comprising a vector, wherein the vector comprises a nucleic acid encoding at least one surface-expressed antibody receptor selected from the group consisting of Igα and Igβ.

- 5 Further provided by the present invention is a hybridoma cell comprising a vector, wherein the vector comprises a nucleic acid encoding at least one surface-expressed antibody receptor selected from the group consisting of Igα and Igβ, wherein the nucleic acid is linked to an inducible functional expression sequence.

- 10 The present invention also provides a method for making a hybridoma cell comprising at least one surface-expressed antibody receptor selected from the group consisting of Igα and Igβ comprising fusing a myeloma cell comprising a vector, wherein the vector comprises a nucleic acid encoding at least one surface-expressed antibody receptor selected from the group consisting of Igα and Igβ, with a B cell to produce a hybridoma cell comprising at least one surface-expressed antibody receptor
15 selected from the group consisting of Igα and Igβ.

The present invention also provides a B cell comprising a vector, wherein the vector comprises a nucleic acid encoding at least one surface-expressed antibody receptor selected from the group consisting of Igα and Igβ.

- 20 Also provided by the present invention is a B cell comprising a vector, wherein the vector comprises a nucleic acid encoding at least one surface-expressed antibody receptor selected from the group consisting of Igα and Igβ, wherein the vector comprises a nucleic acid encoding Igα and Igβ and wherein the vector is integrated into the genome of the B cell.

- 25 The present invention also provides a myeloma cell comprising at least one nucleic acid functionally encoding at least one surface-expressed antibody receptor selected from the group consisting of Igα and Igβ, wherein the nucleic acid encoding the surface-expressed antibody receptor is functionally linked to an inducible expression sequence.

Also provided by the present invention is a myeloma cell comprising at least one nucleic acid functionally encoding at least one mutated surface-expressed antibody receptor selected from the group consisting of Igα and Igβ, wherein the nucleic acid encoding the surface-expressed antibody receptor is functionally linked to an inducible expression sequence.

Also provided by the present invention is a method of making a monoclonal antibody of interest comprising: a) contacting a population of hybridoma cells wherein greater than 15% of the cells in the population express monoclonal antibody that is bound to the cell surface with an antigen linked to a detectable label, wherein the antigen binds to the monoclonal antibody to yield a detectably labeled hybridoma cell; b) isolating the detectably labeled hybridoma cell, thus identifying a hybridoma cell that produces the monoclonal antibody of interest; and c) making the monoclonal antibody of interest from the hybridoma cell.

Also provided by the present invention is a method of making a monoclonal antibody of interest comprising: a) contacting a population of hybridoma cells wherein greater than 15% of the cells in the population express monoclonal antibody that is bound to the cell surface with an antigen, wherein the antigen binds to the monoclonal antibody; b) adding a detectable label to the antigen to yield a detectably labeled hybridoma cell; c) isolating the detectably labeled hybridoma cell, thus identifying a hybridoma cell that produces the monoclonal antibody of interest; and d) making the monoclonal antibody of interest from the hybridoma cell.

Further provided by the present invention is a method of making a monoclonal antibody of interest comprising: a) contacting a hybridoma cell, wherein greater than twenty monoclonal antibody molecules are expressed and bound to the cell surface with an antigen linked to a detectable label, wherein the antigen binds to the monoclonal antibody to yield a detectably labeled hybridoma cell; b) isolating the detectably labeled hybridoma cell, thus identifying a hybridoma cell that produces the monoclonal antibody of interest; and c) making the monoclonal antibody of interest from the hybridoma cell.

Also provided by the present invention is a method of making a monoclonal antibody of interest comprising: a) contacting a hybridoma cell, wherein greater than twenty monoclonal antibody molecules are expressed and bound to the cell surface with an antigen, wherein the antigen binds to the monoclonal antibody; b) adding a
5 detectable label to the antigen to yield a detectably labeled hybridoma cell; c) isolating the detectably labeled hybridoma cell, thus identifying a hybridoma cell that produces the monoclonal antibody of interest; and d) making the monoclonal antibody of interest from the hybridoma cell.

Also provided by the present invention is a method of making a monoclonal
10 antibody of interest comprising: a) contacting a B cell comprising a vector, wherein the vector comprises a nucleic acid encoding at least one surface-expressed antibody receptor selected from the group consisting of Igα and Igβ with an antigen linked to a detectable label, wherein the antigen binds to the monoclonal antibody to yield a detectably labeled B cell; b) isolating the detectably labeled B cell, thus identifying a B
15 cell that produces the monoclonal antibody of interest; and c) making the monoclonal antibody of interest.

The present invention also provides a method of making a monoclonal antibody of interest comprising: a) contacting a B cell comprising a vector, wherein the vector comprises a nucleic acid encoding at least one surface-expressed antibody receptor
20 selected from the group consisting of Igα and Igβ with an antigen; b) adding a detectable label that binds to the antigen to yield a detectably labeled B cell; c) isolating the detectably labeled B cell, thus identifying a B cell that produces the monoclonal antibody of interest; and d) making the monoclonal antibody of interest.

Also provided is a method of making a hybridoma cell that produces a
25 monoclonal antibody that recognizes a selected antigen comprising: a) immunizing a mouse with the antigen; b) fusing a B cell from the immunized mouse with a myeloma cell that comprises at least one nucleic acid functionally encoding at least one surface-expressed antibody receptor selected from the group consisting of Igα and Igβ to produce a monoclonal antibody producing hybridoma cell, wherein the monoclonal

antibody produced by the hybridoma cell is expressed and bound to the cell surface; c) contacting the monoclonal antibody producing hybridoma cell with the antigen, wherein the antigen binds to the monoclonal antibody on the cell surface to produce a detectable hybridoma cell, d) detecting the hybridoma cell and; e) isolating the
5 hybridoma cell, thus making a hybridoma cell that produces a monoclonal antibody that recognizes a specific antigen.

The present invention also provides a transgenic animal comprising B cells comprising a vector, wherein the vector comprises a nucleic acid encoding at least one surface-expressed antibody receptor selected from the group consisting of Ig α and Ig β .
10 The present invention also provides a hematopoietic stem cell comprising a vector, wherein the vector comprises a nucleic acid encoding at least one surface-expressed antibody receptor selected from the group consisting of Ig α and Ig β .

The invention further provides a population of hybridoma cells comprising a vector comprising a nucleic acid encoding Ig α and/or Ig β that expresses monoclonal
15 antibody bound to the cell surface, wherein when the monoclonal antibody is detected by fluorescence, the fluorescence intensity of at least 10% of the cells is at least two fold greater than the fluorescence intensity of a population of hybridoma cells that do not comprise a vector comprising a nucleic acid encoding Ig α and/or Ig β .

The present invention also provides a population of plasma cells comprising a
20 vector comprising a nucleic acid encoding Ig α and/or Ig β that expresses monoclonal antibody bound to the cell surface, wherein when the monoclonal antibody is detected by fluorescence the fluorescence intensity of the population of cells is at least two fold greater than the fluorescence intensity of a population of plasma cells that does not comprise a vector comprising a nucleic acid encoding Ig α and/or Ig β .

25 The present invention also provides a population of hybridoma cells comprising a vector comprising a nucleic acid encoding Ig α and/or Ig β that expresses monoclonal antibody bound to the cell surface, wherein when the monoclonal antibody is detected by fluorescence, the fluorescence intensity of at least 10% of the cells is at least two

fold greater than a the fluorescence intensity of a population of hybridoma cells that do not comprise a vector comprising a nucleic acid encoding Igα and/or Igβ.

Further provided by the present invention a population of plasma cells comprising a vector comprising a nucleic acid encoding Igα and/or Igβ that expresses monoclonal antibody bound to the cell surface, wherein when the monoclonal antibody is detected by fluorescence the fluorescence intensity of at least 10% of the cells is at least two fold greater than the fluorescence intensity of a population of plasma cells that do not comprise a vector comprising a nucleic acid encoding Igα and/or Igβ.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 shows that the surface presentation of antibody such as mIgM in B-cells requires the presence of one or two membrane receptors, Igα and/or Igβ. In B-cells these receptors bind the membrane domain of the membrane form of the heavy chain (mHC, dark gray) portion of an antibody such as mIgM. The small immunoglobulin light chain is drawn in black. In most myelomas the Igα (white) receptor shown here is missing or expressed at very low levels. In some myelomas Igα and/or Igβ may be absent or expressed at low levels. As a result, most hybridomas cannot present significant amounts of mIgM on their surface.

Figure 2 shows the Primer oligonucleotides and PCR amplification of Igα and Igβ cDNAs. (A) The sequences of the sense (S) and antisense (A) primers used to amplify cDNAs encoding Igα and Igβ. These oligos add important restriction endonuclease cloning sites and 5' translation signals (GCCACC) to the receptor sequences. (B) PCR amplification products of the expected sizes in base pairs (bp).

Molecular weight standards flank the PCR products.

Figure 3 shows the sequence of PCR modified mouse Igα cDNA extending from the HindIII to EcoRI cloning sites used to construct Igα expression plasmid p3.1NeoIgα. The Protein sequence is shown above the DNA sequence. The main DNA and protein sequence listings in GenBank for Igα have the accession numbers

NM_007655 and NP_031681, respectively. The cDNA sequence obtained was inconsistent with this original Ig α sequence in a small region, but agrees with the data given by Sakaguchi et al. ("B lymphocyte lineage-restricted expression of mb-1, a gene with CD3-like structural properties" *EMBO J.* 7:3457-64 (1988)) encoding a protein

5 with the sequence encoding six amino acids listed in bold.

Figure 4 shows the sequence of PCR modified mouse Ig β cDNA extending from the HindIII to EcoRI cloning sites used to construct Ig β expression plasmid p3.1NeoIg α . Protein sequence is shown above the DNA sequence. The main sequence listing in GenBank for Ig β has the accession number NM_008339.

10 Figure 5 shows the structure of pcDNA3.1 NeoR vector (Invitrogen, Inc. Life Sciences Division) used to express Ig α receptor protein in transgenic cells.

Figure 6 shows the structure of pcDNA3.1 Zeo vector (Invitrogen, Inc. Life Sciences Division) used to express Ig β receptor protein in transgenic cells.

Figure 7 shows the screening of transfected HGS cell lines for Ig α and Ig β

15 protein levels on western blots. Protein samples (25 μ g of total cell protein in SDS sample buffer) were resolved on a 12% acrylamide gel, transferred to an Immobilon membrane (Millipore), reacted with rabbit polyclonal anti mouse Ig α antibody (A) or Ig β antibody (B), then with donkey anti-rabbit horse antibody radish peroxidase conjugate, and developed following published protocols (Kandasamy et al., 1999).

20 Positions of molecular weight standards (MW Stds) are shown on the left. The range of MWs of the various modified forms of Ig α are shown with a bracket. Other higher bands and bands in control WT lanes appear to be background. Extracts from cell lines are named as follows: SC, spleen cell extract; HGS1, parental hybridoma cell line producing antibodies to GS; and HGS1 α β 1-5 and ECS1 α β 1-3 are the receptor gene

25 transfected cell lines. References of these simplified designations to strain names in laboratory notebooks in the Meagher laboratory at UGA are as follows: HGS1 α β 1, GS-TSC; HGS1 α β 2, GS-TSC-1C; HGS1 α β 3, GS-TE4; HGS1 α β 4, GS-T5D-L; and HGS1 α β 5, GS-TSC-3D. The Ig α and Ig β specific rabbit polyclonal antibodies, anti-MB-1 and SF2B, respectively, were obtained from Dr. Linda Matsuuchi (University of

British Columbia, Vancouver, Canada)(Condon et al., 2000). Antibodies were prepared against synthetic peptides to the cytoplasmic tail of Ig α and the ecto-domain of Ig β .

- Figure 8 shows microscopic examination of increased surface presentation of antibody on hybridoma cells expressing transgenic Ig receptors. All cells shown were treated with FITC labeled goat anti mouse antiserum. All plates were photographed using a 40X Zeiss lens and Hamamatsu digital camera (model C-4742-95) under the same illumination and exposure conditions. Cells treated with an unlabeled antibody have insignificant levels of auto-fluorescence. Cell lines shown are indicated with reference to names in lab notebooks. A & B. Non-transfected HGS1 control cells; D-H. pcDNA3.1-Ig α and -Ig β transfected cells. C HGS1 α β 6, SC-1C#2; D HGS1 α β 7, SC-1CF2; E. HGS1 α β 8, SC-1C#3; F. HGS1 α β 9, SC-3DF3; G. HGS1 α β 10, SC-3DF4; H. HGS1 α β 11, SC-F1. (I) brightest cell enlarged from field of cells in Figure 8A (HGS1 α control hybridoma cells) and indicated by an arrow. (J) Brightest cell enlarged from field of cells in Figure 8C (HGS1 α β 6 receptor transfected hybridoma cells) and indicated by an arrow.

- Figure 9 shows Ig α protein levels in aliquots of HGS1 WT and HGS1 α β cells shown surface presenting antibody in Figure 8. Protein samples were resolved by PAGE and Ig α protein levels measured on Westerns using rabbit antibody to mouse Ig α as in Figure 7A. (A) Ig α western (B) Coomassie stained gel of duplicate samples showing relative level of protein loading. Stained molecular weight standards (MW Stds) are shown on the left in kilo-Daltons (kDa). The range of MWs of the various modified forms of Ig α are shown with a bracket. Extracts from cell lines discussed in Meagher laboratory notebooks are named as follows: WT, HGS1 non transfected control; HGS1 α β 6, SC-1C#2; HGS1 α β 7, SC-1CF2; HGS1 α β 8, SC-1C#3; HGS1 α β 9, SC-3DF3; HGS1 α β 10, SC-3DF4; and HGS1 α β 11, SC-F1. Cell lines showing 100% surface presentation of antibody in Figure 8 are designated plus +, those not presenting as minus, and HGS1 hybridoma WT controls with 1-5% presentation as +/- . Mobility range of Ig α isoforms are shown with a bracket.

Figure 10 shows a western blot examining Ig β expression in aliquots of HGS1 and HGS1 $\alpha\beta$ cells shown in Figure 8. Protein samples were resolved by PAGE and Ig β protein levels measured on Westerns using antibody to Ig β as in Figure 7B. Protein samples are identical aliquots to those shown in Figure 10A and 10B. Mobility range of Ig β isoforms are shown with a bracket.

Figure 11 shows quantification of surface presentation of antibody in Ig α expressing hybridomas. (A) The fluorescence intensity was compared among HGS1 cells and transgenic cells expressing or not expressing detectable levels of Ig α receptor. Mean fluorescence intensity (MFI) was measured for individual cells in a microscopic field from images like those taken in Figure 8, using OpenLab software (Improvision, Inc., Boston, MA). Standard errors in MFI among individual cells in each population are indicated. MFI underestimates the actual intensity differences between control and transgenic cells, because 10% of the brightest cells exceed the dynamic range of the electronic camera and some cells are out of the focal plane, where MFI cannot be accurately measured. n = 50 cells for each of the eight cell populations examined. (B) The frequency of individual cells in each population with 3 times greater intensity than the mean for control HGS1 cells (89 MFI).

Figure 12 shows western blot examining Ig α and Ig β protein levels in transfected myeloma cell lines. Immunodetection of (A) Ig α and (B) Ig β performed as described in Figure 7. Cell lines examined are as follows: SC, spleen cells; Sp2/0 myeloma wild-type (WT) or Sp2/0 derived lines transfected with genes encoding Ig α and Ig β ($\alpha\beta$ 1, M-T4; $\alpha\beta$ 2, M-T4D; $\alpha\beta$ 3, M-T6; $\alpha\beta$ 4, M-T4-3A; $\alpha\beta$ 5, M-T3SC; $\alpha\beta$ 6, M-T1SC; and $\alpha\beta$ 7, M-T4D-6). Mobility ranges of Ig α and Ig β are shown with brackets. Higher molecular weight bands are due to background activity in antibody.

Figure 13 shows fluorescent activated cell sorting (FACS) demonstrating that Ig α expression increases antibody surface presentation eight fold. HGS1 and HGS1 $\alpha\beta$ 10 cells were prepared as shown in Figure 8 and labeled with FITC labeled goat anti-mouse antibody. 200 μ l samples of the following cells were counted (Counts) by FACS using A₅₂₀ emission (FL1). The full scale on the bottom axis represents 4-

logs of fluorescence intensity. A. HGS1 WT control cells. B. HGS1 $\alpha\beta$ 10 cells. C. Mixture of HGS1 and HGS1 $\alpha\beta$ 10 cells.

DETAILED DESCRIPTION OF THE INVENTION

5

The present invention may be understood more readily by reference to the following detailed description of the preferred embodiments of the invention.

Before the present methods are disclosed and described, it is to be understood that this invention is not limited to specific constructs, molecules and methods, as such
10 may of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only and is not intended to be limiting.

It must be noted that, as used in the specification and the appended claims, the singular forms "a," "an," and "the" include plural referents unless the context clearly
15 dictates otherwise. For example, a cell can mean a single cell or more than one cell.

Hybridomas

The present invention provides a population of hybridoma cells wherein greater
20 than 15 % of the cells in the population express monoclonal antibody that is bound to the cell surface. Also provided by this invention is a population of hybridoma cells wherein greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95% or numbers in between, of the cells in the population express monoclonal antibody that is bound to the cell surface.

25 As used herein, "hybridoma" is a cell or a cell line that is produced by fusing an antibody producing cell, e.g. a B cell, and an immortalized cell, e.g. a myeloma cell. As used herein "B cell" means an immature B cell, a mature naïve B cell, a mature activated B cell, a memory B cell, a B lineage lymphocyte, a plasma cell or any other B lineage cell of human origin or from non-human animal sources. The hybridomas of

this invention can be made by fusing a B cell of human origin or from non-human animal sources, with an immortalized cell line using a suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell (*Goding*, "Monoclonal Antibodies: Principles and Practice" Academic Press, (1986) pp. 59-103).

- 5 In order to obtain the B cells for the production of a hybridoma, a mouse or other appropriate host animal, is typically immunized with an immunizing agent or antigen to elicit B cells that produce or are capable of producing antibodies that will specifically bind to the immunizing agent or antigen. Alternatively, the B cells may be immunized *in vitro*. Immortalized cell lines are usually transformed mammalian cells, particularly myeloma cells of rodent, bovine and human origin. Usually, rat or mouse myeloma cell lines are employed. The hybridoma cells may be cultured in a suitable culture medium that preferably contains one or more substances that inhibit the growth or survival of the unfused, immortalized cells. For example, although HAT is not necessary for DISH, typically, if the parental cells lack the enzyme hypoxanthine
- 10 guanine phosphoribosyl transferase (HGPRT or HPRT), the culture medium for the hybridomas typically will include hypoxanthine, aminopterin, and thymidine ("HAT medium"), which substances prevent the growth of HGPRT-deficient cells.

- Preferred immortalized cell lines are those that fuse efficiently, support high level expression of antibody by the selected antibody-producing cells, and are sensitive
- 20 to a medium. The immortalized cell line can be sensitive to HAT medium. More preferred immortalized cell lines are murine myeloma lines, which can be obtained, for instance, from the Salk Institute Cell Distribution Center, San Diego, Calif. and the American Type Culture Collection (ATCC), Rockville, Md. Human myeloma and mouse-human heteromyeloma cell lines also have been described for the production of
- 25 human monoclonal antibodies (*Kozbor*, *J. Immunol.*, 133:3001 (1984) and *Brodeur et al.*, "Monoclonal Antibody Production Techniques and Applications" Marcel Dekker, Inc., New York, (1987) pp. 51-63). For example, the following myeloma cell lines can be obtained from the ATCC: MOPC-31C, RPMI 8226, IM-9, MPC-11, CCL-189, HK-PEG-1, HS-Sultan, A2B5 clone 105, P3X63Ag8.653, Sp2/0-Ag14, Sp2/0-Ag14/SF,

P3X63Ag8U.1, HFN 36.3 HFN 7.1, 45.6.TG1.7, ARH-77, Y3-Ag 1.2.3, SJK-132-20, SJK-287-38 and SJK-237-71.

The hybridoma cells of the present invention can be assayed for surface expression and the culture medium in which the hybridoma cells are cultured can be
 5 assayed for the presence of monoclonal antibodies directed against a desired immunogen by methods known in the art such as ELISA, western blot, FACS, magnetic separation etc. The binding specificity of monoclonal antibodies secreted by the hybridoma cells can be, for example, determined by immunoprecipitation or by an *in vitro* binding assay, such as radioimmunoassay (RIA) or enzyme-linked
 10 immunoabsorbent assay (ELISA). Such techniques and assays are known in the art. The binding affinity of the monoclonal antibody can, for example, be determined by the Scatchard analysis of *Munson et al.*, Anal. Biochem., 107:220 (1980).

After a desired hybridoma cell is identified, either by assaying surface expression or by assaying the culture medium, the selected hybridoma cell can be
 15 grown by standard methods. Suitable culture media for this purpose include, for example, Dulbecco's Modified Eagle's Medium and RPMI-1640 medium. Alternatively, the hybridoma cells may be grown *in vivo* as ascites in a mammal.

As used herein, "a population of hybridoma cells" means a sufficient number of cells such that a percentage of the cells expressing antibody can be determined. The
 20 hybridoma cells of the population can be cells from a pure hybridoma cell line where all of the cells of the line produce only one monoclonal antibody specific for a particular antigen or a mixture of cells wherein multiple monoclonal antibodies are produced. Thus, a population of hybridoma cells can produce more than one monoclonal antibody such that some cells produce a monoclonal antibody that
 25 recognize one antigen and other cells in the population produce monoclonal antibody that recognizes a second antigen and other cells in the population produce a monoclonal antibody that recognizes a third antigen etc.

As used herein, "express" means that the monoclonal antibody can be detected by means standard in the art such as Western blot, ELISA, immunofluorescence,

hemolytic assay, fluorescence activated cell sorting (FACS) as they are currently practiced in the art.

Antibodies are typically proteins which exhibit binding specificity to a specific antigen. Native antibodies are usually heterotetrameric glycoproteins, composed of two identical light (L) chains and two identical heavy (H) chains. Typically, each light chain is linked to a heavy chain by one covalent disulfide bond, while the number of disulfide linkages varies between the heavy chains of different immunoglobulin isotypes. Each heavy and light chain can have regularly spaced intrachain disulfide bridges. Each heavy chain can have at one end a variable domain (V(H)) followed by a number of constant domains. Each light chain can have a variable domain at one end (V(L)) and a constant domain at its other end; the constant domain of the light chain is aligned with the first constant domain of the heavy chain, and the light chain variable domain is aligned with the variable domain of the heavy chain. Particular amino acid residues are believed to form an interface between the light and heavy chain variable domains. The light chains of antibodies from any vertebrate species can be assigned to one of two clearly distinct types, called kappa (κ) and lambda (λ), based on the amino acid sequences of their constant domains. Depending on the amino acid sequence of the constant domain of their heavy chains, immunoglobulins can be assigned to different classes. There currently are five major classes of immunoglobulins: IgA, IgD, IgE, IgG and IgM, and several of these may be further divided into subclasses (isotypes), e.g., IgG-1, IgG-2, IgG-3, and IgG-4; IgA-1 and IgA-2. The present invention provides the presentation of all of the immunoglobulin classes via binding to Ig α and/or Ig β . The heavy chain constant domains that correspond to the different classes of immunoglobulins are called alpha, delta, epsilon, gamma, and mu, respectively.

The Immunoglobulin (Ig) heavy chain genes are typically complex transcription units with multiple poly(A) sites in which changes in the cleavage and polyadenylation machinery can play an important role in B-cell, stage-specific expression. Ig μ heavy chains can be expressed in pre, immature, and mature-B-cells and IgM⁺ plasma cells. The α , ϵ , and γ heavy chains can be expressed in memory and IgA⁺, IgE⁺, and IgG⁺ plasma cells, respectively (Janeway and Travers, 1994). RNA from each of the five classes of Ig heavy chain genes (α , δ , ϵ , γ , μ) can be alternatively processed to produce two types of mRNAs: one encodes the secreted form of the Ig protein and is produced by use of the

promoter-proximal, weak Ig sec (secretory-specific) poly(A) site in plasma cells; the other mRNA encodes the membrane-bound (mb) receptor for antigen on the surface of mature or memory B-cells and can be produced by use of the downstream, strong Ig membrane poly(A) site [Alt, 1980; Rogers, 1980; Rogers, 1981].

- 5 There can be a 2-5-fold change in the transcription rate of the Ig genes in different B-cell stages (Kelly and Perry, 1986). The site of termination can vary in the μ (Galli et al., 1987; Guise et al., 1988; Yuan and Tucker, 1984) but not the γ and α genes (Flaspohler et al., 1995; Flaspohler and Milcarek., 1990; Lebman et al., 1992). RNA processing events can play the major role in determining the ratios of the two forms of
- 10 IgG heavy chain mRNA as first shown in 1985 (Milcarek and Hall, 1985). The crucial role for RNA processing has been further substantiated (See Edwalds-Gilbert and Milcarek, 1995; Edwalds-Gilbert and Milcarek, 1995; Flaspohler et al., 1995; Flaspohler and Milcarek., 1990; Genovese et al., 1991; Genovese and Milcarek, 1990; Hall and Milcarek, 1989; Kobrin et al., 1986; Lassman et al., 1992; Lassman and
- 15 Milcarek, 1992; Matis et al., 1996; Milcarek et al., 1996). See also (Edwalds-Gilbert et al., 1997). Polyadenylation at the weak secretory-specific poly(A) site, which is promoter proximal to the membrane specific poly(A) site, and splicing to the membrane-specific exons at the sub-optimal splice site, in the last secretory-specific exon, can be mutually exclusive events. It has been shown that changes in the cleavage
- 20 and polyadenylation of the precursor RNA tip the balance in plasma cells to the use of the first, weak poly(A) site.

- The term "variable" is used herein to describe certain portions of the variable domains which differ in sequence among antibodies and are used in the binding and specificity of each particular antibody for its particular antigen. However, the
- 25 variability is not usually evenly distributed through the variable domains of antibodies. It is typically concentrated in three segments called complementarity determining regions (CDRs) or hypervariable regions both in the light chain and the heavy chain variable domains. The more highly conserved portions of the variable domains are called the framework (FR). The variable domains of native heavy and light chains can

each comprise four FR regions, largely adopting a β -sheet configuration, connected by three CDRs, which form loops connecting, and in some cases forming part of, the β -sheet structure. The CDRs in each chain can be held together in close proximity by the FR regions and, with the CDRs from the other chain, contribute to the formation of the antigen binding site of antibodies (see *Kabat E. A. et al.*, "Sequences of Proteins of Immunological Interest" National Institutes of Health, Bethesda, Md. (1987)). The constant domains are not typically involved directly in binding an antibody to an antigen, but exhibit various effector functions, such as participation of the antibody in antibody-dependent cellular toxicity.

10 As used herein, "monoclonal antibody" refers to an antibody that is produced by cells that are all derived from a single antibody-producing cell type and has a specific affinity for an antigen. Monoclonal antibodies are obtained from a substantially homogeneous population of antibodies, i.e., the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts. The monoclonal antibodies secreted by the hybridoma cells of the present invention can be isolated or purified from the culture medium or ascites fluid by conventional immunoglobulin purification procedures such as, for example, protein A-Sepharose, hydroxylapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography.

20 Once hybridomas are isolated by the present invention, the antibody coding regions of the hybridomas can be used to make monoclonal antibodies by recombinant DNA methods, such as those described in U.S. Pat. No. 4,816,567 or U.S. Pat. No. 6,331,415. DNA encoding the monoclonal antibodies of the invention can be readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of murine antibodies). The hybridoma cells of the invention serve as a preferred source of such DNA. Once isolated, the DNA may be placed into expression vectors, which are then transfected into host cells such as simian COS cells, Chinese hamster ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin

protein, to obtain the synthesis of monoclonal antibodies in the recombinant host cells.

The DNA also may be modified, for example, by substituting the coding sequence for human heavy and light chain constant domains in place of the homologous murine sequences (U.S. Pat. No. 4,816,567) or by covalently joining to the immunoglobulin

- 5 coding sequence all or part of the coding sequence for a non-immunoglobulin polypeptide. Such a non-immunoglobulin polypeptide can be substituted for the constant domains of an antibody of the invention, or can be substituted for the variable domains of one antigen-combining site of an antibody of the invention to create a chimeric bivalent antibody comprising one antigen-combining site having specificity
10 for one antigen and second antigen-combining site having specificity for a different antigen.

The present invention also provides a hybridoma cell, wherein greater than twenty monoclonal antibody molecules are expressed and bound to the cell surface. Also provided by the present invention is a hybridoma cell, wherein greater than fifty
15 monoclonal antibody molecules are expressed and bound to the cell surface. Further provided by the present invention is a hybridoma cell, wherein greater than one hundred monoclonal antibody molecules are expressed and bound to the cell surface. The present invention also provides a hybridoma cell, wherein greater than two hundred and fifty monoclonal antibody molecules are expressed and bound to the cell surface. Also
20 provided by the present invention is a hybridoma cell, wherein greater than five hundred monoclonal antibody molecules are expressed and bound to the cell surface. Numbers of antibodies in between these numbers are also provided.

- 25 Further provided by the present invention is a population of hybridoma cells wherein greater than 15% of the cells in the population express monoclonal antibody that is bound to the cell surface and wherein greater than twenty monoclonal antibody molecules are expressed and bound to the cell surface of the cells in the population that express monoclonal antibody.

Also provided by the present invention is a population of hybridoma cells wherein greater than 15% of the cells in the population express monoclonal antibody

that is bound to the cell surface and wherein greater than fifty monoclonal antibody molecules are expressed and bound to the cell surface of the cells in the population that express monoclonal antibody.

- Further provided by the present invention is a population of hybridoma cells
- 5 wherein greater than 15% of the cells in the population express monoclonal antibody that is bound to the cell surface and wherein greater than one hundred monoclonal antibody molecules are expressed and bound to the cell surface of the cells in the population that express monoclonal antibody.

- The present invention also provides a population of hybridoma cells wherein
- 10 greater than 15% of the cells in the population express monoclonal antibody that is bound to the cell surface and wherein greater than two hundred and fifty monoclonal antibody molecules are expressed and bound to the cell surface of the cells in the population that express monoclonal antibody.

- Further provided by the present invention is a population of hybridoma cells
- 15 wherein greater than 15% of the cells in the population express monoclonal antibody that is bound to the cell surface and wherein greater than five hundred monoclonal antibody molecules are expressed and bound to the cell surface of the cells in the population that express monoclonal antibody.

- Further provided by the present invention is a population of hybridoma cells
- 20 wherein greater than 25% of the cells in the population express monoclonal antibody that is bound to the cell surface and wherein greater than twenty monoclonal antibody molecules are expressed and bound to the cell surface of the cells in the population that express monoclonal antibody.

- Also provided by the present invention is a population of hybridoma cells
- 25 wherein greater than 25% of the cells in the population express monoclonal antibody that is bound to the cell surface and wherein greater than fifty monoclonal antibody molecules are expressed and bound to the cell surface of the cells in the population that express monoclonal antibody.

Further provided by the present invention is a population of hybridoma cells wherein greater than 25% of the cells in the population express monoclonal antibody that is bound to the cell surface and wherein greater than one hundred monoclonal antibody molecules are expressed and bound to the cell surface of the cells in the
5 population that express monoclonal antibody.

The present invention also provides a population of hybridoma cells wherein greater than 25% of the cells in the population express monoclonal antibody that is bound to the cell surface and wherein greater than two hundred and fifty monoclonal antibody molecules are expressed and bound to the cell surface of the cells in the
10 population that express monoclonal antibody.

Further provided by the present invention is a population of hybridoma cells wherein greater than 25% of the cells in the population express monoclonal antibody that is bound to the cell surface and wherein greater than five hundred monoclonal antibody molecules are expressed and bound to the cell surface of the cells in the
15 population that express monoclonal antibody.

Further provided by the present invention is a population of hybridoma cells wherein greater than 50% of the cells in the population express monoclonal antibody that is bound to the cell surface and wherein greater than twenty monoclonal antibody molecules are expressed and bound to the cell surface of the cells in the population that
20 express monoclonal antibody.

Also provided by the present invention is a population of hybridoma cells wherein greater than 50% of the cells in the population express monoclonal antibody that is bound to the cell surface and wherein greater than fifty monoclonal antibody molecules are expressed and bound to the cell surface of the cells in the population that
25 express monoclonal antibody.

Further provided by the present invention is a population of hybridoma cells wherein greater than 50% of the cells in the population express monoclonal antibody that is bound to the cell surface and wherein greater than one hundred monoclonal

antibody molecules are expressed and bound to the cell surface of the cells in the population that express monoclonal antibody.

The present invention also provides a population of hybridoma cells wherein greater than 50% of the cells in the population express monoclonal antibody that is
5 bound to the cell surface and wherein greater than two hundred and fifty monoclonal antibody molecules are expressed and bound to the cell surface of the cells in the population that express monoclonal antibody.

Further provided by the present invention is a population of hybridoma cells wherein greater than 50% of the cells in the population express monoclonal antibody
10 that is bound to the cell surface and wherein greater than five hundred monoclonal antibody molecules are expressed and bound to the cell surface of the cells in the population that express monoclonal antibody.

Further provided by the present invention is a population of hybridoma cells wherein greater than 75% of the cells in the population express monoclonal antibody
15 that is bound to the cell surface and wherein greater than twenty monoclonal antibody molecules are expressed and bound to the cell surface of the cells in the population that express monoclonal antibody.

Also provided by the present invention is a population of hybridoma cells wherein greater than 75% of the cells in the population express monoclonal antibody
20 that is bound to the cell surface and wherein greater than fifty monoclonal antibody molecules are expressed and bound to the cell surface of the cells in the population that express monoclonal antibody.

Further provided by the present invention is a population of hybridoma cells wherein greater than 75% of the cells in the population express monoclonal antibody
25 that is bound to the cell surface and wherein greater than one hundred monoclonal antibody molecules are expressed and bound to the cell surface of the cells in the population that express monoclonal antibody.

The present invention also provides a population of hybridoma cells wherein greater than 75% of the cells in the population express monoclonal antibody that is

bound to the cell surface and wherein greater than two hundred and fifty monoclonal antibody molecules are expressed and bound to the cell surface of the cells in the population that express monoclonal antibody.

- Further provided by the present invention is a population of hybridoma cells
- 5 wherein greater than 75% of the cells in the population express monoclonal antibody that is bound to the cell surface and wherein greater than five hundred monoclonal antibody molecules are expressed and bound to the cell surface of the cells in the population that express monoclonal antibody.

- Further provided by the present invention is a population of hybridoma cells
- 10 wherein greater than 90% of the cells in the population express monoclonal antibody that is bound to the cell surface and wherein greater than twenty monoclonal antibody molecules are expressed and bound to the cell surface of the cells in the population that express monoclonal antibody.

- Also provided by the present invention is a population of hybridoma cells
- 15 wherein greater than 90% of the cells in the population express monoclonal antibody that is bound to the cell surface and wherein greater than fifty monoclonal antibody molecules are expressed and bound to the cell surface of the cells in the population that express monoclonal antibody.

- Further provided by the present invention is a population of hybridoma cells
- 20 wherein greater than 90% of the cells in the population express monoclonal antibody that is bound to the cell surface and wherein greater than one hundred monoclonal antibody molecules are expressed and bound to the cell surface of the cells in the population that express monoclonal antibody.

- The present invention also provides a population of hybridoma cells wherein
- 25 greater than 90% of the cells in the population express monoclonal antibody that is bound to the cell surface and wherein greater than two hundred and fifty monoclonal antibody molecules are expressed and bound to the cell surface of the cells in the population that express monoclonal antibody.

Further provided by the present invention is a population of hybridoma cells wherein greater than 90% of the cells in the population express monoclonal antibody that is bound to the cell surface and wherein greater than five hundred monoclonal antibody molecules are expressed and bound to the cell surface of the cells in the
5 population that express monoclonal antibody.

Also provided by the present invention is a hybridoma cell, wherein from about 0.001% to about 10% of the total amount of monoclonal antibody produced by the hybridoma cell is expressed and bound to the cell surface.

Any combinations of the above percentages of cells and number of antibodies
10 per cell is also provided as well as numbers in between the specifically listed percentages of cell and number of antibodies per cell.

The total amount of monoclonal antibody produced by a cell can be determined by any of the standard methods in the art, including, but not limited to, Western blot, ELISA, immunofluorescence and FACS. These methods are utilized to measure the
15 amount of antibody secreted into the medium by the cell, the amount of antibody bound to the cell surface and the amount of intracellular antibody present in the cells. One of skill in the art would know how to measure these amounts utilizing the above techniques or others known to the skilled artisan to obtain a total amount of antibody produced by a cell, such that a percentage of the total amount of antibody that is
20 expressed and bound to the cell surface is obtained.

Also provided by the present invention is a hybridoma cell, wherein from about 0.01% to about 10% of the total amount of monoclonal antibody produced by the hybridoma cell is expressed and bound to the cell surface.

Also provided by the present invention is a hybridoma cell, wherein from about
25 10% to about 20% of the total amount of monoclonal antibody produced by the hybridoma cell is expressed and bound to the cell surface.

Also provided by the present invention is a hybridoma cell, wherein from about 20% to about 30% of the total amount of monoclonal antibody produced by the hybridoma cell is expressed and bound to the cell surface.

Also provided by the present invention is a hybridoma cell, wherein from about 30% to about 40% of the total amount of monoclonal antibody produced by the hybridoma cell is expressed and bound to the cell surface.

Also provided by the present invention is a hybridoma cell, wherein from about
5 40% to about 50% of the total amount of monoclonal antibody produced by the hybridoma cell is expressed and bound to the cell surface.

Also provided by the present invention is a hybridoma cell, wherein from about 50% to about 60% of the total amount of monoclonal antibody produced by the hybridoma cell is expressed and bound to the cell surface.

10 Also provided by the present invention is a hybridoma cell, wherein from about 60% to about 70% of the total amount of monoclonal antibody produced by the hybridoma cell is expressed and bound to the cell surface.

Also provided by the present invention is a hybridoma cell, wherein from about 70% to about 80% of the total amount of monoclonal antibody produced by the
15 hybridoma cell is expressed and bound to the cell surface.

Also provided by the present invention is a hybridoma cell, wherein from about 80% to about 90% of the total amount of monoclonal antibody produced by the hybridoma cell is expressed and bound to the cell surface.

Also provided by the present invention is a hybridoma cell, wherein from about
20 90% to about 100% of the total amount of monoclonal antibody produced by the hybridoma cell is expressed and bound to the cell surface.

Further provided by the present invention is a population of hybridoma cells wherein greater than 15% of the hybridoma cells in the population express from about 0.01% to about 10% of the total amount of monoclonal antibody produced by the
25 hybridoma cells on the cell surface. Also provided by this invention is a population of hybridoma cells wherein greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% of the hybridoma cells in the population express from about 0.01% to about 10% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface.

Further provided by the present invention is a population of hybridoma cells wherein greater than 15% of the hybridoma cells in the population express from about 10% to about 20% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface. Also provided by this invention is a population of
5 hybridoma cells wherein greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% of the hybridoma cells in the population express from about 10 % to about 20 % of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface.

Further provided by the present invention is a population of hybridoma cells
10 wherein greater than 15% of the hybridoma cells in the population express from about 20% to about 30% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface. Also provided by this invention is a population of hybridoma cells wherein greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% of the hybridoma cells in the
15 population express from about 20% to about 30% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface.

Further provided by the present invention is a population of hybridoma cells wherein greater than 15% of the hybridoma cells in the population express from about 30% to about 40% of the total amount of monoclonal antibody produced by the
20 hybridoma cells on the cell surface. Also provided by this invention is a population of hybridoma cells wherein greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% of the hybridoma cells in the population express from about 30% to about 40% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface.

25 Further provided by the present invention is a population of hybridoma cells wherein greater than 15% of the hybridoma cells in the population express from about 40% to about 50% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface. Also provided by this invention is a population of hybridoma cells wherein greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%,

60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% of the hybridoma cells in the population express from about 40% to about 50% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface.

Further provided by the present invention is a population of hybridoma cells
5 wherein greater than 15% of the hybridoma cells in the population express from about 50% to about 60% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface. Also provided by this invention is a population of hybridoma cells wherein greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% of the hybridoma cells in the
10 population express from about 50% to about 60% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface.

Further provided by the present invention is a population of hybridoma cells wherein greater than 15% of the hybridoma cells in the population express from about 60% to about 70% of the total amount of monoclonal antibody produced by the
15 hybridoma cells on the cell surface. Also provided by this invention is a population of hybridoma cells wherein greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% of the hybridoma cells in the population express from about 60% to about 70% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface.

Further provided by the present invention is a population of hybridoma cells wherein greater than 15% of the hybridoma cells in the population express from about 70% to about 80% of the total amount of monoclonal antibody produced by the
20 hybridoma cells on the cell surface. Also provided by this invention is a population of hybridoma cells wherein greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% of the hybridoma cells in the
25 population express from about 70% to about 80% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface.

Further provided by the present invention is a population of hybridoma cells wherein greater than 15% of the hybridoma cells in the population express from about

80% to about 90% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface. Also provided by this invention is a population of hybridoma cells wherein greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% of the hybridoma cells in the
5 population express from about 80% to about 90% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface.

Further provided by the present invention is a population of hybridoma cells wherein greater than 15% of the hybridoma cells in the population express from about 90% to about 100% of the total amount of monoclonal antibody produced by the
10 hybridoma cells on the cell surface. Also provided by this invention is a population of hybridoma cells wherein greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% of the hybridoma cells in the population express from about 90% to about 100% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface.

Also provided by the present invention is a population of hybridoma cells wherein greater than 15% of the hybridoma cells in the population express from about 0.01% to about 10% of the total amount of monoclonal antibody produced by the
15 hybridoma cells on the cell surface, and wherein greater than twenty monoclonal antibodies are expressed and bound to the cell surface. Also provided by this invention is a population of hybridoma cells wherein greater than 20%, 25%, 30%, 35%, 40%,
20 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% of the hybridoma cells in the population express from about 0.01% to about 10% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface and wherein greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%,
25 85%, 90%, or 95% of the population are hybridoma cells wherein greater than twenty monoclonal antibodies are expressed and bound to the cell surface.

Also provided by the present invention is a population of hybridoma cells wherein greater than 15% of the hybridoma cells in the population express from about 10% to about 20% of the total amount of monoclonal antibody produced by the

hybridoma cells on the cell surface, and wherein greater than twenty monoclonal antibodies are expressed and bound to the cell surface. Also provided by this invention is a population of hybridoma cells wherein greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% of the hybridoma cells in the population express from about 10% to about 20% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface and wherein greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% of the population are hybridoma cells wherein greater than twenty monoclonal antibodies are expressed and bound to the cell surface.

Also provided by the present invention is a population of hybridoma cells wherein greater than 15% of the hybridoma cells in the population express from about 20% to about 30% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface, and wherein greater than twenty monoclonal antibodies are expressed and bound to the cell surface. Also provided by this invention is a population of hybridoma cells wherein greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% of the hybridoma cells in the population express from about 20% to about 30% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface and wherein greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% of the population are hybridoma cells wherein greater than twenty monoclonal antibodies are expressed and bound to the cell surface.

Also provided by the present invention is a population of hybridoma cells wherein greater than 15% of the hybridoma cells in the population express from about 30% to about 40% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface, and wherein greater than twenty monoclonal antibodies are expressed and bound to the cell surface. Also provided by this invention is a population of hybridoma cells wherein greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% of the hybridoma cells in the population express from about 30% to about 40% of the total amount of

monoclonal antibody produced by the hybridoma cells on the cell surface and wherein greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% of the population are hybridoma cells wherein greater than twenty monoclonal antibodies are expressed and bound to the cell surface.

- 5 Also provided by the present invention is a population of hybridoma cells wherein greater than 15% of the hybridoma cells in the population express from about 40% to about 50% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface, and wherein greater than twenty monoclonal antibodies are expressed and bound to the cell surface. Also provided by this invention
- 10 is a population of hybridoma cells wherein greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% of the hybridoma cells in the population express from about 40% to about 50% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface and wherein greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%,
- 15 85%, 90%, or 95% of the population are hybridoma cells wherein greater than twenty monoclonal antibodies are expressed and bound to the cell surface.

- Also provided by the present invention is a population of hybridoma cells wherein greater than 15% of the hybridoma cells in the population express from about 50% to about 60% of the total amount of monoclonal antibody produced by the
- 20 hybridoma cells on the cell surface, and wherein greater than twenty monoclonal antibodies are expressed and bound to the cell surface. Also provided by this invention is a population of hybridoma cells wherein greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% of the hybridoma cells in the population express from about 50% to about 60% of the total amount of
- 25 monoclonal antibody produced by the hybridoma cells on the cell surface and wherein greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% of the population are hybridoma cells wherein greater than twenty monoclonal antibodies are expressed and bound to the cell surface.

Also provided by the present invention is a population of hybridoma cells wherein greater than 15% of the hybridoma cells in the population express from about 60% to about 70% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface, and wherein greater than twenty monoclonal antibodies are expressed and bound to the cell surface. Also provided by this invention is a population of hybridoma cells wherein greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% of the hybridoma cells in the population express from about 60% to about 70% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface and wherein greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% of the population are hybridoma cells wherein greater than twenty monoclonal antibodies are expressed and bound to the cell surface.

Also provided by the present invention is a population of hybridoma cells wherein greater than 15% of the hybridoma cells in the population express from about 70% to about 80% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface, and wherein greater than twenty monoclonal antibodies are expressed and bound to the cell surface. Also provided by this invention is a population of hybridoma cells wherein greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% of the hybridoma cells in the population express from about 70% to about 80% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface and wherein greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% of the population are hybridoma cells wherein greater than twenty monoclonal antibodies are expressed and bound to the cell surface.

Also provided by the present invention is a population of hybridoma cells wherein greater than 15% of the hybridoma cells in the population express from about 80% to about 90% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface, and wherein greater than twenty monoclonal antibodies are expressed and bound to the cell surface. Also provided by this invention

is a population of hybridoma cells wherein greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% of the hybridoma cells in the population express from about 80% to about 90% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface and wherein
5 greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% of the population are hybridoma cells wherein greater than twenty monoclonal antibodies are expressed and bound to the cell surface.

Also provided by the present invention is a population of hybridoma cells wherein greater than 15% of the hybridoma cells in the population express from about
10 90% to about 100% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface, and wherein greater than twenty monoclonal antibodies are expressed and bound to the cell surface. Also provided by this invention is a population of hybridoma cells wherein greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% of the hybridoma
15 cells in the population express from about 90% to about 100% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface and wherein greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% of the population are hybridoma cells wherein greater than twenty monoclonal antibodies are expressed and bound to the cell surface.

Also provided by the present invention is a population of hybridoma cells wherein greater than 15% of the hybridoma cells in the population express from about 0.01% to about 10% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface, and wherein greater than 15% of the population are hybridoma cells wherein greater than fifty, one hundred, two hundred and fifty or five
20 hundred monoclonal antibodies are expressed and bound to the cell surface. Also
25 provided by this invention is a population of hybridoma cells wherein greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% of the hybridoma cells in the population express from about 0.01% to about 10% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell

surface and wherein greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% of the population are hybridoma cells wherein greater than fifty, one hundred, two hundred and fifty or five hundred monoclonal antibodies are expressed and bound to the cell surface.

- 5 Also provided by the present invention is a population of hybridoma cells wherein greater than 15% of the hybridoma cells in the population express from about 10% to about 20% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface, and wherein greater than 15% of the population are hybridoma cells wherein greater than fifty, one hundred, two hundred and fifty or five
- 10 hundred monoclonal antibodies are expressed and bound to the cell surface. Also provided by this invention is a population of hybridoma cells wherein greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% of the hybridoma cells in the population express from about 10% to about 20% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell
- 15 surface and wherein greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% of the population are hybridoma cells wherein greater than fifty, one hundred, two hundred and fifty or five hundred monoclonal antibodies are expressed and bound to the cell surface.

- Also provided by the present invention is a population of hybridoma cells
- 20 wherein greater than 15% of the hybridoma cells in the population express from about 20% to about 30% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface, and wherein greater than 15% of the population are hybridoma cells wherein greater than fifty, one hundred, two hundred and fifty or five hundred monoclonal antibodies are expressed and bound to the cell surface. Also
- 25 provided by this invention is a population of hybridoma cells wherein greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% of the hybridoma cells in the population express from about 20% to about 30% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface and wherein greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%,

65%, 70%, 75%, 80%, 85%, 90%, or 95% of the population are hybridoma cells wherein greater than fifty, one hundred, two hundred and fifty or five hundred monoclonal antibodies are expressed and bound to the cell surface.

- Also provided by the present invention is a population of hybridoma cells
- 5 wherein greater than 15% of the hybridoma cells in the population express from about 30% to about 40% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface, and wherein greater than 15% of the population are hybridoma cells wherein greater than fifty, one hundred, two hundred and fifty or five hundred monoclonal antibodies are expressed and bound to the cell surface. Also
- 10 provided by this invention is a population of hybridoma cells wherein greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% of the hybridoma cells in the population express from about 30% to about 40% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface and wherein greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%,
- 15 65%, 70%, 75%, 80%, 85%, 90%, or 95% of the population are hybridoma cells wherein greater than fifty, one hundred, two hundred and fifty or five hundred monoclonal antibodies are expressed and bound to the cell surface.

- Also provided by the present invention is a population of hybridoma cells wherein greater than 15% of the hybridoma cells in the population express from about
- 20 40% to about 50% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface, and wherein greater than 15% of the population are hybridoma cells wherein greater than fifty, one hundred, two hundred and fifty or five hundred monoclonal antibodies are expressed and bound to the cell surface. Also provided by this invention is a population of hybridoma cells wherein greater than 20%,
- 25 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% of the hybridoma cells in the population express from about 40% to about 50% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface and wherein greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% of the population are hybridoma cells

wherein greater than fifty, one hundred, two hundred and fifty or five hundred monoclonal antibodies are expressed and bound to the cell surface.

- Also provided by the present invention is a population of hybridoma cells wherein greater than 15% of the hybridoma cells in the population express from about
- 5 50% to about 60% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface, and wherein greater than 15% of the population are hybridoma cells wherein greater than fifty, one hundred, two hundred and fifty or five hundred monoclonal antibodies are expressed and bound to the cell surface. Also provided by this invention is a population of hybridoma cells wherein greater than 20%,
- 10 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% of the hybridoma cells in the population express from about 50% to about 60% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface and wherein greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% of the population are hybridoma cells
- 15 wherein greater than fifty, one hundred, two hundred and fifty or five hundred monoclonal antibodies are expressed and bound to the cell surface.

- Also provided by the present invention is a population of hybridoma cells wherein greater than 15% of the hybridoma cells in the population express from about 60% to about 70% of the total amount of monoclonal antibody produced by the
- 20 hybridoma cells on the cell surface, and wherein greater than 15% of the population are hybridoma cells wherein greater than fifty, one hundred, two hundred and fifty or five hundred monoclonal antibodies are expressed and bound to the cell surface. Also provided by this invention is a population of hybridoma cells wherein greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%,
- 25 75%, 80%, 85%, 90%, or 95% of the hybridoma cells in the population express from about 60% to about 70% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface and wherein greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% of the population are hybridoma cells wherein greater than fifty, one hundred, two hundred

and fifty or five hundred monoclonal antibodies are expressed and bound to the cell surface.

- Also provided by the present invention is a population of hybridoma cells wherein greater than 15% of the hybridoma cells in the population express from about
- 5 70% to about 80% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface, and wherein greater than 15% of the population are hybridoma cells wherein greater than fifty, one hundred, two hundred and fifty or five hundred monoclonal antibodies are expressed and bound to the cell surface. Also provided by this invention is a population of hybridoma cells wherein greater than 20%,
- 10 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% of the hybridoma cells in the population express from about 70% to about 80% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface and wherein greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% of the population are hybridoma cells
- 15 wherein greater than fifty, one hundred, two hundred and fifty or five hundred monoclonal antibodies are expressed and bound to the cell surface.

- Also provided by the present invention is a population of hybridoma cells wherein greater than 15% of the hybridoma cells in the population express from about 80% to about 90% of the total amount of monoclonal antibody produced by the
- 20 hybridoma cells on the cell surface, and wherein greater than 15% of the population are hybridoma cells wherein greater than fifty, one hundred, two hundred and fifty or five hundred monoclonal antibodies are expressed and bound to the cell surface. Also provided by this invention is a population of hybridoma cells wherein greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or
- 25 95% of the hybridoma cells in the population express from about 80% to about 90% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface and wherein greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% of the population are hybridoma cells

wherein greater than fifty, one hundred, two hundred and fifty or five hundred monoclonal antibodies are expressed and bound to the cell surface.

- Also provided by the present invention is a population of hybridoma cells wherein greater than 15% of the hybridoma cells in the population express from about
- 5 90% to about 100% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface, and wherein greater than 15% of the population are hybridoma cells wherein greater than fifty, one hundred, two hundred and fifty or five hundred monoclonal antibodies are expressed and bound to the cell surface. Also provided by this invention is a population of hybridoma cells wherein greater than 20%,
- 10 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% of the hybridoma cells in the population express from about 90% to about 100% of the total amount of monoclonal antibody produced by the hybridoma cells on the cell surface and wherein greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% of the population are hybridoma cells
- 15 wherein greater than fifty, one hundred, two hundred and fifty or five hundred monoclonal antibodies are expressed and bound to the cell surface.

- Also provided by this invention is a population of hybridoma cells comprising monoclonal antibodies bound to the cell surface having about two, three, four, five, six, seven, eight, nine, ten times more antibody on the cell surface than control cells, i.e.
- 20 standard hybridoma cells.

- Further provided by this invention is a population of hybridoma cells comprising a vector comprising a nucleic acid encoding Ig α and/or Ig β that expresses monoclonal antibody bound to the cell surface, wherein when the monoclonal antibody is detected by fluorescence, the fluorescence intensity of the population of cells is at
- 25 least two fold greater than the fluorescence intensity of a population of hybridoma cells that do not comprise a vector comprising a nucleic acid encoding Ig α and/or Ig β .

The invention also provides a population of hybridoma cells comprising a vector comprising a nucleic acid encoding Ig α and/or Ig β that expresses monoclonal antibody bound to the cell surface, wherein when the monoclonal antibody is detected

by fluorescence, the fluorescence intensity of the population of cells is at least two fold, three fold, four fold, five fold, six fold, seven fold, eight fold, nine fold, ten fold, fifteen fold, twenty fold, thirty fold, forty fold, fifty fold, sixty fold, seventy fold, eighty fold, ninety fold, one hundred fold, two hundred and fifty fold, five hundred fold or one thousand fold greater than the fluorescence intensity of a population of hybridoma cells that do not comprise a vector comprising a nucleic acid encoding Ig α and/or Ig β . The fold increase in fluorescence intensity can also be any amount in between the fold increases listed above. The fold increase in fluorescence intensity can be measured by methods standard in the art and is described herein in the Examples.

The population of hybridoma cells utilized to measure fluorescence intensity can be between 25 and 500 cells. Therefore, the population can be about 25, 50, 75, 100, 125, 150, 175, 200, 225, 250, 275, 300, 325, 350, 375, 400, 425, 450, 475, 500 cells or any number of cells in between these values.

The hybridomas in the population can comprise a vector comprising a nucleic acid encoding Ig α .

Further provided by the present invention is a population of hybridoma cells comprising a vector comprising a nucleic acid encoding Ig α and/or Ig β that expresses monoclonal antibody bound to the cell surface, wherein when the monoclonal antibody is detected by fluorescence, the fluorescence intensity of at least 10% of the cells is at least two fold greater than the fluorescence intensity of a population of hybridoma cells that do not comprise a vector comprising a nucleic acid encoding Ig α and/or Ig β .

The present invention also provides a population of hybridoma cells comprising a vector comprising a nucleic acid encoding Ig α and/or Ig β that expresses monoclonal antibody bound to the cell surface, wherein when the monoclonal antibody is detected by fluorescence, the fluorescence intensity of at least 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100% or any percentage in between of the cells is at least two fold greater than the fluorescence intensity of a population of hybridoma cells that do not comprise a vector comprising a nucleic acid encoding Ig α and/or Ig β .

The present invention also provides a population of hybridoma cells comprising a vector comprising a nucleic acid encoding Ig α and/or Ig β that expresses monoclonal antibody bound to the cell surface, wherein when the monoclonal antibody is detected by fluorescence, the fluorescence intensity of at least 10%, 20%, 30%, 40%, 50%, 60%,
5 70%, 80%, 90%, 100% or any percentage in between of the cells is at least two fold, three fold, five fold, six fold, seven fold, eight fold, nine fold, ten fold, twenty fold, thirty fold, forty fold, fifty fold, sixty fold, seventy fold, eight fold, ninety fold, one hundred fold, two hundred and fifty fold, five hundred fold, one thousand fold or any amount in between, greater than the fluorescence intensity of a population of
10 hybridoma cells that do not comprise a vector comprising a nucleic acid encoding Ig α and/or Ig β .

Further provided by the present invention is a hybridoma cell comprising a vector, wherein the vector comprises a nucleic acid encoding at least one surface-expressed antibody receptor selected from the group consisting of Ig α and Ig β .
15 Therefore, the hybridoma can comprise a vector comprising a nucleic acid encoding Ig α , or the hybridoma can comprise a vector comprising a nucleic acid encoding Ig β , or the hybridoma can comprise a vector comprising a nucleic acid encoding both Ig α and a nucleic acid encoding Ig β . The nucleic acids encoding Ig α or Ig β can be present in a single vector or in multiple vectors. For example, the hybridoma can comprise a vector
20 comprising a nucleic acid encoding Ig α and a vector comprising a nucleic acid encoding Ig β or a vector comprising the nucleic acid sequences for both Ig α and Ig β . Any of the vectors can be integrated into the genome of the cell or carried extrachromosomally to allow transient expression of Ig α and/or Ig β .

As used herein, the term "nucleic acid" refers to single-or multiple stranded
25 molecules which may be DNA or RNA, or any combination thereof, including modifications to those nucleic acids. The nucleic acid may represent a coding strand or its complement, or any combination thereof. Nucleic acids may be identical in sequence to the sequences which are naturally occurring for the Ig α and Ig β receptors discussed herein or may include alternative codons which encode the same amino acid

as that which is found in the naturally occurring sequence. Also contemplated are nucleic acid sequences encoding Ig α and Ig β that contain deletions, substitutions, mutations and combinations thereof as long as the nucleic acid encodes a functional receptor. These nucleic acids can also be modified from their typical structure. Such
 5 modifications include, but are not limited to, methylated nucleic acids, the substitution of a non-bridging oxygen on the phosphate residue with either a sulfur (yielding phosphorothioate deoxynucleotides), selenium (yielding phosphorselenoate deoxynucleotides), or methyl groups (yielding methylphosphonate deoxynucleotides).

The nucleic acid sequence for the Ig α receptor is available from GenBank via
 10 Accession No. NM_007655 and the polypeptide encoded by this nucleic acid sequence is available from GenBank via Accession Number NP_031681. The nucleic acid sequence encoding Ig α that was utilized in the Examples described herein differs from the original Ig α receptor available from GenBank via Accession No. NM_007655, but is consistent with the sequence provided by Sakaguchi et al. This sequence is provided
 15 in Figure 3.

A nucleic acid molecule encoding Ig α and a nucleic acid encoding Ig β can be isolated from the organism in which it is normally found. For example, a genomic DNA or cDNA library can be constructed and screened for the presence of the nucleic acid of interest. Methods of constructing and screening such libraries are well known
 20 in the art and kits for performing the construction and screening steps are commercially available (for example, Stratagene Cloning Systems, La Jolla, CA). Once isolated, the nucleic acid can be directly cloned into an appropriate vector, or if necessary, be modified to facilitate the subsequent cloning steps. Such modification steps are routine, an example of which is the addition of oligonucleotide linkers which contain restriction
 25 sites to the termini of the nucleic acid. General methods are set forth in *Sambrook et al.*, "Molecular Cloning, a Laboratory Manual," Cold Spring Harbor Laboratory Press (1989).

Once the nucleic acid sequence of the desired Ig α and/or Ig β is obtained, the sequence encoding specific amino acids can be modified or changed at any particular

amino acid position by techniques well known in the art. For example, PCR primers can be designed which span the amino acid position or positions and which can substitute any amino acid for another amino acid. Then a nucleic acid can be amplified and inserted into the wild-type receptor coding sequence in order to obtain any of a

- 5 number of possible combinations of amino acids at any position of the receptors. Alternatively, one skilled in the art can introduce specific mutations at any point in a particular nucleic acid sequence through techniques for point mutagenesis. General methods are set forth in *Smith, M.* "In vitro mutagenesis" *Ann. Rev. Gen.*, 19:423-462 (1985) and *Zoller, M.J.* "New molecular biology methods for protein engineering" *Curr. Opin. Struct. Biol.*, 1:605-610 (1991). Techniques such as these can be used to alter the coding sequence without altering the amino acid sequence that is encoded.

Another example of a method of obtaining a DNA molecule encoding Ig α is to synthesize a recombinant DNA molecule which encodes Ig α . A nucleic acid encoding Ig β can also be obtained in this manner. For example, oligonucleotide synthesis

- 15 procedures are routine in the art and oligonucleotides coding for a particular protein region are readily obtainable through automated DNA synthesis. A nucleic acid for one strand of a double-stranded molecule can be synthesized and hybridized to its complementary strand. One can design these oligonucleotides such that the resulting double-stranded molecule has either internal restriction sites or appropriate 5' or 3' overhangs at the termini for cloning into an appropriate vector. Double-stranded molecules coding for relatively large proteins can readily be synthesized by first constructing several different double-stranded molecules that code for particular regions of the protein, followed by ligating these DNA molecules together. For example, *Cunningham, et al.*, "Receptor and Antibody Epitopes in Human Growth Hormone Identified by Homolog-Scanning Mutagenesis," *Science*, 243:1330-1336 (1989), have constructed a synthetic gene encoding the human growth hormone gene by first constructing overlapping and complementary synthetic oligonucleotides and ligating these fragments together. See also, *Ferretti, et al.*, *Proc. Nat. Acad. Sci.* 82:599-603 (1986), wherein synthesis of a 1057 base pair synthetic bovine rhodopsin gene from

synthetic oligonucleotides is disclosed. By constructing a nucleic acid in this manner, one skilled in the art can readily obtain any particular Ig α or Ig β with desired amino acids at any particular position or positions within the Ig α or Ig β . See also, U.S. Patent No. 5,503,995 which describes an enzyme template reaction method of making

- 5 synthetic genes. Techniques such as this are routine in the art and are well documented. These nucleic acids or fragments of a nucleic acid encoding Ig α or Ig β can then be expressed *in vivo* or *in vitro* as discussed below. Similarly, nucleic acids or fragments of a nucleic acid encoding can be expressed *in vivo* or *in vitro*.

- Once a nucleic acid encoding Ig α or a region of that nucleic acid, is constructed,
- 10 modified, or isolated, that nucleic acid can then be cloned into an appropriate vector, which can direct the *in vivo* or *in vitro* synthesis of that wild-type and/or modified Ig α receptor protein. Also, once a nucleic acid encoding Ig β or a region of that nucleic acid, is constructed, modified, or isolated, that nucleic acid can then be cloned into an appropriate vector, which can direct the *in vivo* or *in vitro* synthesis of that wild-type
- 15 and/or modified Ig β receptor protein. The vector is contemplated to have the necessary functional elements that direct and regulate transcription of the inserted gene, or nucleic acid. These functional elements include, but are not limited to, a promoter, regions upstream or downstream of the promoter, such as enhancers that may regulate the transcriptional activity of the promoter, an origin of replication, appropriate restriction
- 20 sites to facilitate cloning of inserts adjacent to the promoter, antibiotic resistance genes or other markers which can serve to select for cells containing the vector or the vector containing the insert, RNA splice junctions, a transcription termination region, or any other region which may serve to facilitate the expression of the inserted gene or hybrid gene. (See generally, *Sambrook et al.*). One could also transfect DNA or RNA
- 25 encoding an extra copy of Ig α and/or Ig β directly into the cell in the absence of additional functional elements, e.g. as naked DNA or RNA, as long as the Ig α and/or Ig β resulted in increased expression of the receptor.

The vector comprising a nucleic acid encoding Ig α and Ig β of the present invention can be any vector suitable for expression of a nucleic acid in a eukaryotic cell

as are known in the art. For example, pcDNA3.1 NeoR vector or pcDNA 3.1 Zeo can be utilized (Invitrogen, Inc. Life Sciences Division). Other vectors include, but are not limited to, a two vector inducible system from Invitrogen (pIND and pVgRXXR plasmids), a two vector inducible system from Clontech (pTet-ON or pTet-Off and pTRE2 plasmids), single plasmids for constitutive expression from Promega (pCI or pSI plasmids), a two vector inducible system from Stratagene (pCMVLacI and pOPRSVI/MCS plasmids), single plasmid inducible systems from Stratagene (pERV3 or pEGSH plasmids) and single retroviral inducible systems from Stratagene (pCFB-EGSH or pFB-ERV retroviral vectors). The vector can also be a viral vector such as an adenoviral vector, an adeno-associated viral vector, a retroviral vector, a lentiviral vector, a pseudotyped retroviral vector, or a pox virus vector, such as a vaccinia virus vector.

The present invention also provides a hybridoma cell comprising a vector, wherein the vector comprises a nucleic acid encoding at least one mutated surface-expressed antibody receptor selected from the group consisting of Ig α and Ig β . Thus, the present invention provides a hybridoma cell comprising a vector that comprises a nucleic acid encoding a mutant Ig α receptor, a hybridoma cell comprising a vector that comprises a nucleic acid encoding a mutant Ig β receptor, and a hybridoma cell that comprises a vector that comprises both a nucleic acid encoding mutant Ig α receptor and a nucleic acid encoding a mutant Ig β receptor.

The mutant Ig α and Ig β receptors include non-signalling receptors with altered cytoplasmic domains. An example of such a mutant receptor is an Ig α receptor comprising mutations at amino acid residues 176, 182, 193 and 204. The mutated Ig α receptors of the present invention also include a mutated Ig α that comprises one or more mutations selected from the group consisting of Y176F, Y182F, Y193F and Y204F. Further provided by the present invention is a mutated Ig α receptor that comprises a deletion of amino acid residues 176-220. Another example of a mutant surface expressed antibody receptor is a mutated Ig β receptor comprising mutations at amino acid residues 190 and 206. The mutated Ig β receptors of the present invention

also include a mutated $\text{Ig}\beta$ receptor comprising one or more mutations selected from the group consisting of Y190F and Y206F. The mutations described herein for the $\text{Ig}\alpha$ and the $\text{Ig}\beta$ are designed such that the receptors retain the ability to bind antibodies, but are unable to act as signaling receptors. One of skill in the art would know how to

5 manipulate the nucleic acids encoding $\text{Ig}\alpha$ and $\text{Ig}\beta$ as described above as well as other techniques known in the art to obtain the mutant receptors described herein as well as other mutant receptors.

Also provided by the present invention is a hybridoma cell comprising a vector, wherein the vector comprises a nucleic acid encoding at least one chimeric surface-

10 expressed antibody receptor selected from the group consisting of $\text{Ig}\alpha$ and $\text{Ig}\beta$. As used throughout this application, "chimeric" means that the cell surface receptor can comprise a sequence derived from a receptor sequence of one species, e.g. human, and a receptor sequence derived from another species. For example, a chimeric $\text{Ig}\alpha$ can comprise a human $\text{Ig}\alpha$ extracellular domain and a mouse $\text{Ig}\alpha$ transmembrane domain and a

15 mouse intracellular (cytoplasmic) domain or a chimeric $\text{Ig}\alpha$ can comprise a human $\text{Ig}\alpha$ extracellular domain and a human transmembrane domain and a mouse intracellular domain. Similarly, a chimeric $\text{Ig}\beta$ can comprise a human $\text{Ig}\beta$ extracellular domain and a mouse $\text{Ig}\beta$ transmembrane and mouse intracellular domain or a chimeric $\text{Ig}\beta$ can comprise a human $\text{Ig}\beta$ extracellular domain and a human $\text{Ig}\beta$ transmembrane and a

20 mouse intracellular domain. Receptor sequences from other species such as chicken, dog, rabbit, rat, gerbil and hamster can also be utilized to make the chimeric receptors of the present invention. Other examples of chimeric receptors include, but are not limited to a chimeric $\text{Ig}\alpha$ or $\text{Ig}\beta$ receptor comprising a rabbit N-terminal extracellular domain, a mouse transmembrane domain and a mouse C-terminal intracellular domain; a chimeric

25 receptor comprising a chicken N-terminal domain, a mouse transmembrane domain and a mouse C-terminal domain; a chimeric receptor comprising a mouse N-terminal extracellular domain, a chicken transmembrane domain and a chicken C-terminal intracellular signaling domain; a chimeric receptor comprising a mouse N-terminal extracellular domain, a rabbit transmembrane domain and a human C-terminal

intracellular signaling domain; a chimeric receptor comprising a mouse N-terminal extracellular domain a human transmembrane domain and a mutant C-terminal intracellular non-signaling domain from mouse.”

5 The chimeric receptors of this invention also include chimeric receptors comprising a sequence derived from Ig α or Ig β and another non-related sequence. For example, the present invention contemplates a chimeric Ig α receptor comprising an extracellular domain from a non-related protein, such as CD8 or any other protein with an extracellular domain and a transmembrane Ig α domain and an intracellular Ig α domain.

10 The present invention further provides a hybridoma cell comprising a vector, wherein the vector comprises a nucleic acid encoding at least one surface-expressed antibody receptor selected from the group consisting of Ig α and Ig β , wherein the nucleic acid is linked to an inducible functional expression sequence.

15 All of the sequences encoding Ig α and Ig β can be functionally linked to an expression sequence. The expression sequences can include a promoter, an enhancer, a silencer and necessary information processing sites, such as ribosome binding sites, RNA splice sites, polyadenylation sites and transcriptional terminator sequences. The promoters utilized can be constitutive promoters or inducible promoters.

20 The inducible expression systems that can be used for the compositions and methods of the present invention include the IPTG based regulatory system, a tetracycline based regulatory system, CID based regulatory system, an ecdysone based regulatory system, and an estrogen-based regulatory system. Burcin et al., “A Regulatory System for Target Gene Expression” *Frontiers in Bioscience*, 3:1-7 (1998) describes these systems in detail and is incorporated herein in its entirety for the
25 purposes of describing these inducible expression systems. Another inducible system that can be utilized is the cre-lox system (See Lakso “Targeted oncogene activation by site-specific recombination in transgenic mice.” *Proc. Natl Acad Sci USA* 89: 6861-65 (1992); Orban et al., “Tissue and site-specific DNA recombination in transgenic mice” *Proc Natl Acad Sci USA* 90: 6861-65 (1992); Gu et al., “Deletion of a DNA polymerase

beta gene segment in T cells using cell type-specific gene targeting" *Science* 265: 103-106 (1994)). The nucleic acids of the present invention can also be under the control of an inducible metallothionine promoter (See Cox and Maness "Neurite extension and protein tyrosine phosphorylation elicited by inducible expression of the v-src oncogene in a PC12 cell line" *Exp Cell Res* 195: 423-31 (1991)).

In addition to comprising vectors comprising a nucleic acid encoding at least one surface expressed antibody receptor selected from the group consisting of Ig α and Ig β , the hybridomas of the present invention can also comprise a vector comprising a nucleic acid encoding U1A, an enzyme involved in inhibiting the expression of the secretory form of immunoglobulin M (See Philips et al., "Regulation of nuclear poly (A) addition controls the expression of immunoglobulin M secretory mRNA, *EMBO* 22:6443-6452 (2001)).

All of the hybridoma cells described in this application can be utilized in the methods described herein to make a monoclonal antibody of interest.

Methods of Making Hybridomas

Also provided by the present invention is a method for making a hybridoma cell comprising at least one surface-expressed antibody receptor selected from the group consisting of Ig α and Ig β comprising fusing a myeloma cell comprising a vector, wherein the vector comprises a nucleic acid encoding at least one surface-expressed antibody receptor selected from the group consisting of Ig α and/or Ig β , with a B cell to produce a hybridoma cell comprising at least one surface-expressed antibody receptor selected from the group consisting of Ig α and Ig β .

In the methods of making the hybridoma cells of the present invention, the vector can integrate into the genome of the cell. The vector may also be carried extrachromosomally in the cell, thus allowing transient expression of Ig α and Ig β . In the methods of making the hybridomas of the present invention, the nucleic acids encoding Ig α and Ig β can be functionally linked to an inducible expression sequence.

Inducible expression systems are discussed above.

1 The myeloma cells of the present invention can comprise at least one nucleic acid functionally encoding at least one surface-expressed antibody receptor selected from the group consisting of Ig α and Ig β , wherein the nucleic acid encoding the surface-expressed antibody receptor is functionally linked to an inducible expression sequence. The myeloma cells of the present invention can also comprise a nucleic acid encoding a mutated Ig α receptor and/or a mutated Ig β receptor. An example of such a mutant receptor is an Ig α receptor comprising mutations at amino acid residues 176, 182, 193 and 204. The mutated Ig α receptors of the present invention also include a mutated Ig α that comprises one or more mutations selected from the group consisting of Y176F, Y182F, Y193F and Y204F. Further provided by the present invention is a myeloma cell comprising a nucleic acid encoding a mutated Ig α receptor that comprises a deletion of amino acid residues 176-220. Another example of a mutant surface expressed antibody receptor is a mutated Ig β receptor comprising mutations at amino acid residues 190 and 206. The mutated Ig β receptors of the present invention also include a mutated Ig β receptor comprising one or more mutations selected from the group consisting of Y190F and Y206F. The myeloma cell of the present invention can be a myeloma cell comprising a vector that encodes a mutant Ig α receptor, a myeloma cell comprising a vector that comprises a mutant Ig β receptor, or a myeloma cell that comprises a mutant Ig α receptor and a mutant Ig β receptor.

The myeloma cell utilized in the methods of the present invention can also comprise a nucleic acid encoding a chimeric Ig α receptor and/or a chimeric Ig β receptor.

25

B cells

The present invention also provides a B cell comprising a vector, wherein the vector comprises a nucleic acid encoding at least one surface-expressed antibody receptor selected from the group consisting of Ig α and Ig β . Therefore, the B cells of

this invention can comprise a vector comprising a nucleic acid encoding Ig α , or a vector comprising a nucleic acid encoding Ig β , or a vector comprising a nucleic acid encoding both Ig α and Ig β .

- The B cells of the present invention can also comprise a vector comprising a
- 5 nucleic acid encoding a mutated Ig α receptor and/or a mutated Ig β receptor. An example of such a mutant receptor is an Ig α receptor comprising mutations at amino acid residues 176, 182, 193 and 204. The mutated Ig α receptors of the present invention also include a mutated Ig α that comprises one or more mutations selected from the group consisting of Y176F, Y182F, Y193F and Y204F. Further provided by
- 10 the present invention is a B cell comprising a vector comprising a nucleic acid encoding a mutated Ig α receptor that comprises a deletion of amino acid residues 176-220. Another example of a mutant surface expressed antibody receptor is a mutated Ig β receptor comprising mutations at amino acid residues 190 and 206. The mutated Ig β receptors of the present invention also include a mutated Ig β receptor comprising one or
- 15 more mutations selected from the group consisting of Y190F and Y206F. The B cell of the present invention can be a B cell comprising a vector that encodes a mutant Ig α receptor, a B cell comprising a vector that encodes a mutant Ig β receptor, or a B cell that comprises a vector encoding a mutant Ig α receptor and a mutant Ig β receptor. The B cells of the present invention can also comprise a nucleic acid encoding a chimeric
- 20 Ig α receptor and/or a chimeric Ig β receptor.

- The present invention also provides a B cell comprising a vector, wherein the vector comprises a nucleic acid encoding Ig α and Ig β , wherein the nucleic acid encoding Ig α and Ig β is functionally linked to an inducible expression sequence and wherein the nucleic acid encoding Ig α and Ig β is integrated into the genome of the cell.
- 25 Such a B cell can be obtained from the transgenic animals described herein.

The present invention also provides a method of making a B cell comprising a vector, wherein the vector comprises a nucleic acid encoding at least one surface-expressed antibody receptor selected from the group consisting of Ig α and Ig β comprising the steps of transfecting a B cell with a vector comprising at least one

nucleic acid functionally encoding at least one surface-expressed antibody receptor selected from the group consisting of Ig α and Ig β , wherein the nucleic acid encoding the surface-expressed antibody receptor is functionally linked to an expression sequence. In the methods of making the B cells of the present invention, the nucleic acids encoding Ig α and Ig β can be functionally linked to an inducible expression sequence. In the methods of making the B cells of the present invention, the vector comprising at least one nucleic acid functionally encoding at least one surface-expressed antibody receptor selected from the group consisting of Ig α and Ig β can be integrated into the genome of the B cell. Alternatively, the vector does not integrate into the genome of the cell and the vector is carried extrachromosomally to allow transient expression of Ig α and/or Ig β .

The vectors of the present invention can be transfected into cells using any technique known in the art. For example, lipofectamine transfection, microinjection, electroporation, liposomal delivery and particle gun bombardment can all be utilized to effect vector delivery to cells.

In order to transfect B cells removed from an animal's spleen, B cells can be propagated for 24-48 hours in culture so that they divide. A retroviral vector comprising a nucleic acid encoding comprising at least one nucleic acid functionally encoding at least one surface-expressed antibody receptor selected from the group consisting of Ig α and Ig β can then be transfected into the cells. In order to promote proliferation, cytokines can be added to the culture. Cytokines that can be utilized in these methods include, but are not limited to, IL-2, IL-4, IL-5, IL-6, IFN- γ , and/or TGF- β . B cells that produce antibodies bound to their cell surface can then be detected by methods known in the art and described herein, such as FACS, cell panning, ELISA etc. Alternatively, a viral vector such as an adenoviral vector, a lentiviral vector, an adeno-associated vector, a vaccinia virus vector, a pseudotyped retroviral vector can be utilized to transfect B cells. One of skill in the art would know how to test B cells for their ability to be transfected by a vector and select the vector that is most suitable for the introduction of nucleic acids into these cells. One of skill in the art can also

engineer the B cells of the present invention to produce a cell surface receptor that would be recognized by a particular vector. For example, the B cells of the present invention could be engineered such that a cell-surface receptor for adenovirus is present on the cell surface of the B cells in order to facilitate entry of the adenoviral vector into the B cells. Alternatively, a viral vector comprising a ligand that binds to a receptor (other than $Ig\alpha$ and $Ig\beta$) that has been introduced into the B cell can also be utilized to effect vector transfer. The presence of these cell surface receptors can be controlled by an inducible expression system described herein or elsewhere in the art, such that after transfection, expression of the cell surface receptor necessary for viral entry is no longer induced and thus the receptor is no longer expressed on the cell surface of the B cell. These B cells can then be fused to an immortalized cell, such as a naturally occurring myeloma or a genetically altered myeloma cell, such as those provided herein to make a hybridoma cell line expressing a monoclonal antibody of interest. Alternatively the amino acid sequence of the antibody or desired portion thereof, such as a variable region made by such B cell, or the nucleic acid (cDNA) that codes for such antibody or portion thereof, may be determined or isolated. Such DNA sequence may then be incorporated into a vector which directs the expression and secretion of such antibody and such vector transfected into a host cell. such as a myeloma or other appropriate immortal cell. Techniques for determining transfecting and expressing such antibody sequences are described in U.S. Pat. No. 5,627,052 and U.S. Pat. No. 6,331,415.

The B cells of the present invention can further comprise a detectable label.

The present invention also provides an immortalized B cell made by transfecting the B cell with a nucleic acid encoding telomerase. (See Bunk "Immortalizing Human Cells, *The Scientist* 14:19 (2000)). Immortalized B cells can also be made be inactivating pRB/p16 (INK4a) in addition to enhanced telomerase expression (See Kiyono et al., 1998; Dickson et al., 2000). Furthermore, immortalized cells can be made by overexpressing c-myc and simian virus 40 large T antigen (Greenberg et al., 1999; Kim et al., 2001). Immortalized B cells can also be made by

overexpressing Cyclin D1 and inactivating p 53 (See Opitz et al., 2001) or by overexpressing SV40 large T antigen alone (Russo et al., 1998). Other methods of immortalizing B cells include overexpressing *ras* genes and overexpressing human papillomavirus 16E6 and E7 genes (See Coursen et al., 1997). Another combination of
5 genes that can be utilized is hTERT, sv40 large T oncoprotein and an onco-allele of H-ras.

For some applications it may be desirable to generate B cells that are capable of expressing one or more Ig receptors and are also immortal. One means of achieving this is to use embryos derived from an animal that is transgenic for one or more
10 immortalizing genes. One such animal is an IMMORTOMOUSE® mouse, commercially available through Charles River Laboratories. Such mice have a temperature sensitive SV40 T antigen gene in most cells. Those of ordinary skill in the art will recognize that immortal B cells may also be obtained by using transgenic animals that carry additional genes known to immortalize cells as described above such
15 as hTERT or H-ras.

In addition to comprising vectors comprising a nucleic acid encoding at least one surface expressed antibody receptor selected from the group consisting of Igα and Igβ, the B cells of the present invention can also comprise a vector comprising a nucleic acid encoding U1A, an enzyme involved in inhibiting the expression of the secretory
20 form of immunoglobulin M (See Philips et al., "Regulation of nuclear poly (A) addition controls the expression of immunoglobulin M secretory mRNA, *EMBO* 22:6443-6452 (2001).

All of the B cells comprising vectors described herein can be fused to a myeloma cell or other immortal cell line to make a hybridoma cell. The resulting
25 hybridoma cell can be utilized in the methods of making a monoclonal antibody of interest described herein.

Plasma cells

There are approximately 10^8 B cells in a typical mouse spleen. About 99% of such B cells surface present antibody. However 1%, or 10% of such cells are plasma cells and typically surface present only trace amounts of antibody. Plasma cells are known to produce immunoglobulin that is highly specific and of strong affinity for particular target antigens. This invention provides populations of plasma cells that surface present adequate immunoglobulin to enable high throughput fluorescence activated cell sorting technology to be used to determine whether single cells produce immunoglobulin that react with target antigens. Such plasma cells may be obtained from any animal such as the transgenic animals provided herein and can produce fully human immunoglobulin, if isolated from a transgenic animal that expresses a nucleic acid coding for human antibodies in its B cells.

As stated above, plasma cells make very specific, high affinity antibodies to target antigens. Therefore, it is desirable to isolate plasma cells from among a larger population of B cells prior to sorting the plasma cells to identify cells that produce desired antibodies. The marker SYNDECAN-1 is expressed to a higher degree on plasma cells than on other B cells. In addition, plasma cells do not express IGD or B220, whereas other B cells do express both markers. Commercial antibodies for SYNDECAN, IGD and B220 are available and the three markers may be used by those of ordinary skill in the art to segregate plasma cells from among B cell populations by methods known in the art. Plasma cells may also be separated from other cells by density-based centrifugation where the fraction containing plasma cells is collected using an elutriator. Alternatively, a purified plasma cell population may be achieved using separation/purification columns such as those that utilizing magnetic beads.

The present invention provides a population of plasma cells wherein greater than 5 % of the cells in the population express monoclonal antibody that is bound to the cell surface. Also provided by this invention is a population of plasma cells wherein greater than 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%,

75%, 80%, 85%, 90%, or 95% of the cells in the population express monoclonal antibody that is bound to the cell surface.

The present invention also provides a plasma cell, wherein greater than twenty monoclonal antibody molecules are expressed and bound to the cell surface. Also provided by the present invention is a plasma cell, wherein greater than fifty monoclonal antibody molecules are expressed and bound to the cell surface. Further provided by the present invention is a plasma cell, wherein greater than one hundred monoclonal antibody molecules are expressed and bound to the cell surface. The present invention also provides a plasma cell, wherein greater than two hundred and fifty monoclonal antibody molecules are expressed and bound to the cell surface. Also provided by the present invention is a plasma cell, wherein greater than five hundred monoclonal antibody molecules are expressed and bound to the cell surface.

Further provided by this invention is a population of plasma cells comprising a vector comprising a nucleic acid encoding Ig α and/or Ig β that expresses monoclonal antibody bound to the cell surface, wherein when the monoclonal antibody is detected by fluorescence, the fluorescence intensity of the population of cells is at least two fold greater than the fluorescence intensity of a population of plasma cells that do not comprise a vector comprising a nucleic acid encoding Ig α and/or Ig β .

The invention also provides a population of plasma cells comprising a vector comprising a nucleic acid encoding Ig α and/or Ig β that expresses monoclonal antibody bound to the cell surface, wherein when the monoclonal antibody is detected by fluorescence, the fluorescence intensity of the population of cells is at least two fold, three fold, four fold, five fold, six fold, seven fold, eight fold, nine fold, ten fold, fifteen fold, twenty fold, thirty fold, forty fold, fifty fold, sixty fold, seventy fold, eighty fold, ninety fold, one hundred fold, two hundred and fifty fold, five hundred fold, one hundred fold, two hundred and fifty fold, five hundred fold or one thousand fold greater than the fluorescence intensity of a population of plasma cells that do not comprise a vector comprising a nucleic acid encoding Ig α and/or Ig β . The fold increase in fluorescence intensity can also be any amount in between the fold increases listed

above. The fold increase in fluorescence intensity can be measured by methods standard in the art and is described herein in the Examples.

The population of plasma cells utilized to measure fluorescence intensity can be between 25 and 500 cells. Therefore, the population can be about 25, 50, 75, 100, 125,
 5 150, 175, 200, 225, 250, 275, 300, 325, 350, 375, 400, 425, 450, 475, 500 cells or any number of cells in between these values.

The plasma cells in the population can comprise a vector comprising a nucleic acid encoding Ig α .

Further provided by the present invention is a population of plasma cells
 10 comprising a vector comprising a nucleic acid encoding Ig α and/or Ig β that expresses monoclonal antibody bound to the cell surface, wherein when the monoclonal antibody is detected by fluorescence, the fluorescence intensity of at least 10% of the cells is at least two fold greater than the fluorescence intensity of a population of plasma cells that do not comprise a vector comprising a nucleic acid encoding Ig α and/or Ig β .

The present invention also provides a population of plasma cells comprising a
 15 vector comprising a nucleic acid encoding Ig α and/or Ig β that expresses monoclonal antibody bound to the cell surface, wherein when the monoclonal antibody is detected by fluorescence, the fluorescence intensity of at least 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100% or any percentage in between of the cells is at least two fold greater
 20 than the fluorescence intensity of a population of plasma cells that do not comprise a vector comprising a nucleic acid encoding Ig α and/or Ig β .

The present invention also provides a population of plasma cells comprising a
 vector comprising a nucleic acid encoding Ig α and/or Ig β that expresses monoclonal
 25 antibody bound to the cell surface, wherein when the monoclonal antibody is detected by fluorescence, the fluorescence intensity of at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100% or any percentage in between of the cells is at least two fold, three fold, five fold, six fold, seven fold, eight fold, nine fold, ten fold, twenty fold, thirty fold, forty fold, fifty fold, sixty fold, seventy fold, eight fold, ninety fold, one hundred fold, two hundred and fifty fold, five hundred fold, one thousand fold or any

amount in between, greater than the fluorescence intensity of a population of plasma cells that do not comprise a vector comprising a nucleic acid encoding Igα and/or Igβ.

All of the populations of plasma cells described herein can be utilized in the methods of making antibodies provide by the present invention such that the plasma
 5 cells of the present invention can be contacted with an antigen/antigens in order to identify monoclonal antibody producing cells which can be isolated and subsequently produced.

Myeloma cells

10 The present invention also provides a myeloma cell that comprises at least one nucleic acid functionally encoding at least one surface-expressed antibody receptor selected from the group consisting of Igα and Igβ, wherein the nucleic acid encoding the surface-expressed antibody receptor is functionally linked to an inducible
 15 expression sequence.

Further provided by the present invention is a myeloma cell that comprises at least one nucleic acid functionally encoding at least one mutated surface-expressed antibody receptor selected from the group consisting of Igα and Igβ, wherein the nucleic acid encoding the surface-expressed antibody receptor is functionally linked to
 20 an inducible expression sequence.

Also provided by the present invention is a myeloma cell that comprises a nucleic acid functionally encoding a mutated Igα receptor having a deletion of amino acid residues 176- 220, wherein the nucleic acid encoding the surface-expressed antibody receptor is functionally linked to an inducible expression sequence.

25 Further provided by this invention is a myeloma cell that comprises a nucleic acid functionally encoding a mutated Igα receptor having one or more mutations selected from the group consisting of: Y176F, Y182F, Y193F, Y204F, wherein the nucleic acid encoding the surface-expressed antibody receptor is functionally linked to an inducible expression sequence.

The present invention further provides a myeloma cell that comprises a nucleic acid functionally encoding a mutated Ig β receptor having one or more mutations selected from the group consisting of: Y190F and Y206F, wherein the nucleic acid encoding the surface-expressed antibody receptor is functionally linked to an inducible expression sequence.

Any of the myeloma cells comprising vectors provided by the present invention can be fused to a B cell, including a B cell comprising a vector of the present invention, to make a hybridoma cell. The resulting hybridoma cell can then be used in the methods of making monoclonal antibodies described herein.

10 The present invention also provides a method of making a myeloma cell that comprises at least one nucleic acid functionally encoding at least one surface-expressed antibody receptor selected from the group consisting of Ig α and Ig β , wherein the nucleic acid encoding the surface-expressed antibody receptor is functionally linked to an inducible expression sequence comprising the steps of transfecting a myeloma cell
15 with at least one nucleic acid functionally encoding at least one surface-expressed antibody receptor selected from the group consisting of Ig α and Ig β , wherein the nucleic acid encoding the surface-expressed antibody receptor is functionally linked to an inducible expression sequence.

The present invention also provides a method of making a myeloma cell that
20 comprises at least one nucleic acid functionally encoding at least one mutated surface-expressed antibody receptor selected from the group consisting of Ig α and Ig β , wherein the nucleic acid encoding the surface-expressed antibody receptor is functionally linked to an inducible expression sequence comprising transfecting a myeloma cell with at least one nucleic acid functionally encoding at least one mutated surface-expressed
25 antibody receptor selected from the group consisting of Ig α and Ig β , wherein the nucleic acid encoding the surface-expressed antibody receptor is functionally linked to an inducible expression sequence.

Also provided by the present invention is a method of making a myeloma cell that comprises a nucleic acid functionally encoding a mutated Ig α receptor having one

or more mutations selected from the group consisting of: Y176F, Y182F, Y193F, Y204F, wherein the nucleic acid encoding the surface-expressed antibody receptor is functionally linked to an inducible expression sequence comprising transfecting a myeloma cell with a nucleic acid functionally encoding a mutated Igα receptor having
5 one or more mutations selected from the group consisting of: Y176F, Y182F, Y193F, Y204F, wherein the nucleic acid encoding the surface-expressed antibody receptor is functionally linked to an inducible expression sequence.

The present invention provides a method of making a myeloma cell that comprises a nucleic acid functionally encoding a mutated Igβ receptor having one or
10 more mutations selected from the group consisting of: Y190F and Y206F, wherein the nucleic acid encoding the surface-expressed antibody receptor is functionally linked to an inducible expression sequence comprising transfecting a myeloma cell with a nucleic acid functionally encoding a mutated Igβ receptor having one or more mutations selected from the group consisting of: Y190F and Y206F, wherein the
15 nucleic acid encoding the surface-expressed antibody receptor is functionally linked to an inducible expression sequence.

The present invention provides a method of making a myeloma cell that comprises a nucleic acid functionally encoding a mutated Igα receptor having a deletion of amino acid residues 176- 220, wherein the nucleic acid encoding the surface-
20 expressed antibody receptor is functionally linked to an inducible expression sequence comprising transfecting a myeloma cell with a nucleic acid functionally encoding a mutated Igα receptor having a deletion of amino acid residues 176- 220, wherein the nucleic acid encoding the surface-expressed antibody receptor is functionally linked to an inducible expression sequence.

In addition to comprising vectors comprising a nucleic acid encoding at least
25 one surface expressed antibody receptor selected from the group consisting of Igα and Igβ, the myeloma cells of the present invention can also comprise a vector comprising a nucleic acid encoding U1A, an enzyme involved in inhibiting the expression of the secretory form of immunoglobulin M (See Philips et al., "Regulation of nuclear poly

(A) addition controls the expression of immunoglobulin M secretory mRNA, *EMBO* 22:6443-6452 (2001).

The present invention also provides a method of screening myeloma cells or other other immortal cells for the presence of Ig α and Ig β on their surface. If a myeloma cell or an immortal cell naturally expressing Ig α and Ig β is identified, this cell can be fused to B cells to produce hybridoma cells expressing monoclonal antibodies on their cell surface. If a myeloma cell or an immortal cell naturally expressing Ig α is identified, this cell can be fused to the B cells of the present invention to produce hybridoma cells expressing monoclonal antibodies on their cell surface.

Alternatively, this cell can be transfected with a vector comprising a nucleic acid encoding Ig β and then fused to a B cell in order to produce hybridoma cells expressing monoclonal antibodies on their cell surface. If a myeloma cell or an immortal cell naturally expressing Ig β is identified, this cell can be fused to the B cells of the present invention to produce hybridoma cells expressing monoclonal antibodies on their cell surface. Alternatively, this cell can be transfected with a vector comprising a nucleic acid encoding Ig α and then fused to a B cell in order to produce hybridoma cells expressing monoclonal antibodies on their cell surface.

Myeloma cells can also be screened to determine which of the myeloma cells is a suitable fusion partner for making hybridomas. One of skill in the art would know how to test myelomas for desirable fusion characteristics either before or after screening for the presence of Ig α and/or Ig β in order to determine which ones are best suited for fusion with B cells. Alternatively, once a myeloma cell or immortalized cell is deemed to be suitable for fusion, this myeloma cell can be transfected with Ig α and/or Ig β prior to fusion with a B cell. In addition, since HAT selection is not required, the investigator's choice of the cell to be used as a fusion partner for B cells in a given protocol is greatly expanded. Using DISH, myelomas or other candidate fusion partners can be identified that are more cell sparing or offer other advantages over standard myelomas in use today.

Methods of Making Monoclonal Antibodies

The present invention provides method of making a monoclonal antibody of interest comprising: a) contacting a population of hybridoma cells wherein greater than
5 15 % of the cells in the population express monoclonal antibody that is bound to the cell surface with an antigen linked to a detectable label, wherein the antigen binds to the monoclonal antibody to yield a detectably labeled hybridoma cell; b) isolating the detectably labeled hybridoma cell, thus identifying a hybridoma cell that produces the monoclonal antibody of interest; c) making the monoclonal antibody of interest from
10 the hybridoma cell.

Also provided is a method of making a monoclonal antibody of interest comprising: a) contacting a population of hybridoma cells wherein greater than 15 % of the cells in the population express monoclonal antibody that is bound to the cell surface with an antigen, wherein the antigen binds to the monoclonal antibody ; b) adding a
15 detectable label to the antigen to yield a detectably labeled hybridoma cell; c) isolating the detectably labeled hybridoma cell, thus identifying a hybridoma cell that produces the monoclonal antibody of interest; d) making the monoclonal antibody of interest from the hybridoma cell.

In the methods of making a monoclonal antibody described herein, conditions
20 whereby an antigen/antibody complex can form as well as assays for the detection of the formation of an antigen/antibody complex and quantitation of the detected protein are standard in the art. Such assays can include, but are not limited to, Western blotting, immunoprecipitation, immunofluorescence, immunocytochemistry, immunohistochemistry, fluorescence activated cell sorting (FACS), fluorescence in situ
25 hybridization (FISH), immunomagnetic assays, ELISA, ELISPOT (Coligan et al., eds., Current Protocols in Immunology, Wiley, New York (1995)), agglutination assays, flocculation assays, cell panning, magnetic separation etc., as are well known to those of skill in the art.

The antigen of this invention can be bound to a substrate (e.g., beads, tubes, slides, plates, nitrocellulose sheets, etc.) or conjugated with a detectable label (moiety) or both bound and conjugated. The detectable moieties contemplated for the present invention can include, but are not limited to, an immunofluorescence moiety (e.g., fluorescein, rhodamine), a radioactive moiety (e.g., ^{32}P , ^{125}I , ^{35}S), an enzyme moiety (e.g., horseradish peroxidase, alkaline phosphatase), a colloidal gold moiety, a dye and a biotin moiety. Such conjugation techniques are standard in the art (see, e.g., Harlow and Lane, "Antibodies, A Laboratory Manual" Cold Spring Harbor Publications, New York, (1988); Yang et al., *Nature* 382: 319-324 (1996)). Labels can be coupled either directly or indirectly to the antigens. One example of indirect coupling is by use of a spacer moiety. These spacer moieties, in turn, can be either insoluble or soluble (Diener et al., *Science* 231: 148 (1986)).

There are many different labels and methods of labeling known to those of ordinary skill in the art. Examples of the types of labels which can be used in the present invention include enzymes, radioisotopes, fluorescent compounds, chemiluminescent compounds, microspheres dyes and bioluminescent compounds. Furthermore, the binding of these labels to the antigens required for practice of the invention can be done using standard techniques common to those of ordinary skill in the art.

Since the populations of hybridomas described herein can produce more than one monoclonal antibody, the present invention provides for methods of making monoclonal antibodies, wherein the population of cells is contacted with more than one antigen. Once each antigen binds a monoclonal antibody of interest, each one can be detected by a separate label, thus identifying more than one monoclonal antibody of interest in a population of cells. For example, the population can be contacted with three antigens, wherein each antigen is labeled either directly or indirectly with a different fluorescent label. The monoclonal antibody producing cells can be detected and the three different monoclonal antibody producing cells can be distinguished based on the differences in fluorescence associated with the different labels. Therefore, the

present invention allows the isolation and production of more than one monoclonal antibody from a population of cells. This same approach can be applied to the isolation and production of multiple monoclonal antibodies from the B cells of this invention, including the isolation and production of monoclonal antibodies from the plasma cells of this invention.

In the above methods of making a monoclonal antibody of interest, a population of hybridoma cells wherein greater than 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% can be contacted with the antigen.

A method of making a monoclonal antibody of interest comprising: a) contacting a hybridoma cell, wherein greater than twenty monoclonal antibody molecules are expressed and bound to the cell surface with an antigen linked to a detectable label, wherein the antigen binds to the monoclonal antibody to yield a detectably labeled hybridoma cell; b) isolating the detectably labeled hybridoma cell, thus identifying a hybridoma cell that produces the monoclonal antibody of interest; c) making the monoclonal antibody of interest from the hybridoma cell.

A method of making a monoclonal antibody of interest comprising: a) contacting a hybridoma cell, wherein greater than twenty monoclonal antibody molecules are expressed and bound to the cell surface with an antigen, wherein the antigen binds to the monoclonal antibody; b) adding a detectable label to the antigen to yield a detectably labeled hybridoma cell; c) isolating the detectably labeled hybridoma cell, thus identifying a hybridoma cell that produces the monoclonal antibody of interest; d) making the monoclonal antibody of interest from the hybridoma cell.

In the above methods of making a monoclonal antibody of interest, a hybridoma cell wherein greater than fifty, one hundred, two hundred and fifty or five hundred monoclonal antibody molecules are expressed and bound to the cell surface can be contacted with the antigen.

In all of the methods of making a monoclonal antibody of interest, the hybridoma cells can comprise a vector comprising a nucleic acid encoding at least one surface-expressed antibody receptor selected from the group consisting of Ig α and Ig β . The nucleic acid encoding at least one surface-expressed antibody receptor selected from the group consisting of Ig α and Ig β can be a mutated Ig α and/or mutated Ig β , a chimeric Ig α and/or chimeric Ig β as described above.

The present invention also provides a method of making a monoclonal antibody of interest comprising: a) contacting a B cell comprising a vector, wherein the vector comprises a nucleic acid encoding at least one surface-expressed antibody receptor selected from the group consisting of Ig α and Ig β with an antigen linked to a detectable label, wherein the antigen binds to the monoclonal antibody to yield a detectably labeled B cell; b) isolating the detectably labeled B cell, thus identifying a B cell that produces the monoclonal antibody of interest; and c) making the monoclonal antibody of interest.

The present invention also provides a method of making a monoclonal antibody of interest comprising: a) contacting a B cell comprising a vector, wherein the vector comprises a nucleic acid encoding at least one surface-expressed antibody receptor selected from the group consisting of Ig α and Ig β with an antigen; b) add a detectable label that binds to the antigen to yield a detectably labeled B cell; c) isolating the detectably labeled B cell, thus identifying a B cell that produces the monoclonal antibody of interest; and d) making the monoclonal antibody of interest.

The invention further provides a method of making a monoclonal antibody of interest comprising: a) contacting a B cell comprising a vector, wherein the vector comprises a nucleic acid encoding at least one surface-expressed antibody receptor selected from the group consisting of Ig α and Ig β with an antigen linked to a detectable label, wherein the antigen binds to the monoclonal antibody to yield a detectably labeled B cell; b) isolating the detectably labeled B cell, thus identifying a B cell that produces the monoclonal antibody of interest; c) determining the amino acid sequence of the variable region of the monoclonal antibody; and d) making the monoclonal

antibody of interest. The amino acid sequence of the variable region of the monoclonal antibody can be determined by obtaining RNA from an antibody producing cell, such as a B cell, constructing a cDNA, amplifying the cDNA by utilizing primers corresponding to a DNA sequence in the variable region of the immunoglobulin chain, determining the nucleotide sequence and translating the nucleotide sequence in order to obtain an amino acid sequence for the variable region of the monoclonal antibody. For the purposes of determining the amino acid sequence of the variable region of a monoclonal antibody, please see, U.S. Patent 5,627,052 which is hereby incorporated in its entirety by this reference.

The invention further provides a method of making a monoclonal antibody of interest comprising: a) contacting a B cell comprising a vector, wherein the vector comprises a nucleic acid encoding at least one surface-expressed antibody receptor selected from the group consisting of Ig α and Ig β with an antigen linked to a detectable label, wherein the antigen binds to the monoclonal antibody to yield a detectably labeled B cell; b) isolating the detectably labeled B cell, thus identifying a B cell that produces the monoclonal antibody of interest; c) obtaining a nucleic acid encoding the variable region of the monoclonal antibody and d) making the monoclonal antibody of interest. A nucleic acid encoding the variable region of the monoclonal antibody can be obtained by isolating DNA from an antibody producing cell, such as a B cell. Once DNA is isolated, the DNA sequences encoding the rearranged variable regions, including the complementarity determining regions are amplified by PCR and the resulting amplification product sequenced. For the purposes of obtaining a nucleic acid encoding the variable region of a monoclonal antibody, please see U.S. Patent 5, 627, 052 which is hereby incorporated in its entirety by this reference.

In all of the methods of making a monoclonal antibody of interest from B cells, the B cells can comprise a vector comprising a nucleic acid encoding at least one surface-expressed antibody receptor selected from the group consisting of Ig α and Ig β . The nucleic acid encoding at least one surface-expressed antibody receptor selected

from the group consisting of Ig α and Ig β can be a mutated Ig α and/or mutated Ig β , a chimeric Ig α and/or chimeric Ig β .

The present invention also provides a method of making a hybridoma cell that produces a monoclonal antibody that recognizes a selected antigen comprising: a) immunizing a mouse with the antigen; b) fusing a B cell from the immunized mouse with a myeloma cell that comprises at least one nucleic acid functionally encoding at least one surface-expressed antibody receptor selected from the group consisting of Ig α and Ig β to produce a monoclonal antibody producing hybridoma cell, wherein the monoclonal antibody produced by the hybridoma cell is expressed and bound to the cell surface; c) contacting the monoclonal antibody producing hybridoma cell with the antigen, wherein the antigen binds to the monoclonal antibody on the cell surface to produce a detectable hybridoma cell, and d) isolating the detectable hybridoma cell, thus making a hybridoma cell that produces a monoclonal antibody that recognizes a specific antigen. In this method, the antigen can be directly labeled to yield a detectably labeled hybridoma cell.

As used herein, an "antigen" can be a peptide, a polypeptide, a recombinant polypeptide, a carbohydrate, a nucleic acid, a lipid, a fragment of a polypeptide, such as a C-terminal fragment or an N-terminal fragment, an organic compound, a synthetic compound, a naturally occurring compound derived from bacterial, plant, animal, protist or fungal source. The antigen can also comprise the binding site of a cell surface receptor such that monoclonal antibodies against that particular site can be made to target a cell surface receptor.

A method of making a hybridoma cell that produces a monoclonal antibody that recognizes a selected antigen comprising: a) contacting a B cell comprising a vector, wherein the vector comprises a nucleic acid encoding at least one surface-expressed antibody receptor selected from the group consisting of Ig α and Ig β with an antigen wherein the antigen binds to the monoclonal antibody to yield a detectable B cell; b) isolating the detectable B cell, thus identifying a B cell that produces the monoclonal antibody of interest and; c) fusing the B cell that produces the monoclonal antibody of

interest to a myeloma cell to produce a hybridoma cell that produces a monoclonal antibody that recognizes a selected antigen. In this method, the antigen can be directly labeled to yield a detectably labeled B cell.

- Also provided by this invention is a transgenic animal comprising B cells
- 5 comprising a vector, wherein the vector comprises a nucleic acid encoding at least one surface-expressed antibody receptor selected from the group consisting of Ig α and Ig β functionally linked to expression sequences, including but not limited to a promoter, intronic sequences and poly-adenylation signal sequences. The B cells comprising a vector can comprise at least one mutated surface-expressed antibody receptor selected
- 10 from the group consisting of Ig α and Ig β . The B cells comprising a vector can comprise at least one chimeric surface-expressed antibody receptor selected from the group consisting of Ig α and Ig β . The B cells comprising a vector can comprise a mutated Ig α receptor comprising one or more mutations selected from the group consisting of: Y176F, Y182F, Y193F, Y204F. The B cells comprising a vector can
- 15 comprise a mutated Ig β receptor comprising one or more mutations selected from the group consisting of: Y190F and Y206F.

- The transgenic animals of this invention can be made by methods known in the art. For the purposes of generating a transgenic animal, screening the transgenic animal for the presence of a transgene and other methodology regarding transgenic animals,
- 20 please see U.S. Patent No. 6,111,166 which is incorporated by this reference in its entirety. For example, the transgenic animals of this invention can be made by a) injecting a transgene comprising a nucleic acid encoding Ig α functionally linked to an expression sequence and/or a transgene comprising a nucleic acid encoding Ig β functionally linked to an expression sequence into an embryo and b) allowing the
- 25 embryo to develop into an animal. This can further comprise crossing the animal with a second animal to produce a third animal. B cells comprising a transgene, wherein the transgene comprises a nucleic acid encoding at least one surface-expressed antibody receptor selected from the group consisting of Ig α and Ig β can be isolated from the

transgenic animal of this invention. The transgenic animals of the present invention include, but are not limited to, mouse, rat, rabbit, guinea pig.

In the transgenic animals of the present invention, the transgene can be expressed in immature B cells, mature naive B cells, mature activated B cells, memory
5 B cells, B lineage lymphocytes and/or plasma cells. Therefore, the expression sequences can be selected such that expression of the transgene is directed to B cells, but not exclusively to B cells. The expression sequence can direct expression to one, more than one or all of the following types of B cells: immature B cells, mature naive B cells, mature activated B cells, memory B cells, B lineage lymphocytes and plasma
10 cells.

In the transgenic animals of the present invention, expression of the transgene can be controlled by an inducible promoter. The transgenic animal of this invention can utilize an inducible expression system such as the *cre-lox*, metallothioneine, or tetracycline-regulated transactivator system. Using the example of the *cre-lox* system,
15 the genes of interest (Ig α and Ig β) are inserted onto a plasmid or suitable viral vector containing a stop codon flanked by locus of crossing over (*loxP*) sites which comprise two 13 base pair inverted repeats separated by an 8 base pair spacer region. This cassette is under control of a specific promoter such as the immunoglobulin kappa, immunoglobulin lambda, CD19, CD45R/B220, CD81 (TAPA-1), or CD138 (syndecan-
20 1) promoter. The genes of interest are inserted in the plasmid on the opposite side of the *loxP*-stop-*loxP* region from the cell specific promoter. In another plasmid, *cre*-recombinase is inserted next to a promoter whose expression may be controlled (proI). Each plasmid is micro-injected into the pronuclei of separate embryos and the embryos implanted into a pseudopregnant female. Additionally, the plasmids may be used to
25 transform embryonic stem cells from a suitable animal. The latter will thereafter be combined with blastocysts from the same or similar non-human animal and re-implanted into pseudopregnant foster mothers to generate chimeric animals comprising the plasmid comprising the transgene. Further methods of generating transgenic animals well known in the art, such as lipofectin or viral transfection of embryonic stem

cells or pre-implantation embryos, may also be used. Alternatively, mice bearing a proI-*cre* transgene may include already established mice such as the interferon inducible 'Mx-Cre' mouse by Kuhn et al. (see below).

Transgenic animals are mated and the resulting F1 animals are screened for the gene via PCR and/or Southern blot analysis. After homozygosity for the transgene is established, animals possessing the proI-*cre* sequence are then mated with animals with an intact pro-*loxP*-stop-*loxP*-Ig α -Ig β . The resulting F1 animals are then screened for individuals possessing both transgenes by PCR and/or Southern blot analysis. In the case of the 'Mx-Cre' cre recombinase transgene, expression of the pro-*loxP*-stop-*loxP*-Ig α -Ig β transgene is achieved by initiating expression of the *cre* recombinase such as through the injection of type-1 interferon (IFN) as is the case with the 'Mx-Cre' cre recombinase transgene. The *cre*-recombinase will then initiate a recombination event targeted at the *loxP* sites by binding at the inverted repeats of one *lox* site and forming a synapse with the second site. *Cre*-recombinase will then cleave the DNA in the spacer region and initiate strand exchange between the synapsed *loxP* sites. This will result in the deletion of the stop codon and transcription from the promoter through the Ig α and Ig β genes. A similar method is detailed by M. Lasko et al., in "Targeted oncogene activation by site-specific recombination in transgenic mice," PNAS, 89, 6232-6236, July 1992, and is included herein in its entirety. Though this is only one method of using the cre/*lox* system similar results may be achieved by inserting the Ig α and Ig β genes onto a plasmid or viral vector in reverse orientation to the promoter and between *loxP* sites in opposite orientation (pro-*loxP*-Ig β -Ig α -*loxP*). In this scenario, once a recombination event is initiated the genes may reverse orientation (pro-Ig α -Ig β) allowing transcription. An example of this is documented in M. Mitsou et al., "Memory B-cell persistence is independent of persisting immunizing antigen," *Nature* 407, 636-642, Oct. 5, 2000 and included herein in its entirety. The use of Mx-Cre transgenic mouse and type-1 IFN as an inducer was published by R. Kuhn et al., "Inducible gene targeting in mice," *Science*, 269(5229): 1427-1429, Sept. 8, 1995, and is included herein by reference in its entirety.

In another approach, B cells from animals with an intact pro-*IoxP*-stop-*IoxP*-*Igα*-*Igβ* will be treated *in vitro* with a cell permeable Cre recombinase protein such as that described by Jo *et al.*, "Epigenetic regulation of gene structure and function with a cell-permeable Cre recombinase," *Nature Biotechnology*, 19: 929-933, 2001, and is

5 included herein by reference in its entirety.

The transgenic animals of the present invention can also utilize a tetracycline system where the genes of interest (*Igα* and/or *Igβ*) are inserted into a plasmid or viral vector adjacent to a tetracycline-responsive promoter (TRE). In another plasmid, tetracycline-controlled transactivator (rtTA) is inserted next to a promoter that can

10 direct expression to B cells or a constitutive promoter. As with the cre-lox system transgenic animals may be made by micro-injection of pronuclei or stem cell transformation. The resulting F1 animals are screened for the gene. Animals possessing the pro-rtTA sequence are bred to homozygosity and then mated with animals with an intact TRE- *Igα*-*Igβ*. The resulting F1 animals are then screened for

15 individuals possessing both transgenes. Expression of the transgene is achieved by injecting tetracycline or a suitable derivative such as doxycycline. The dox will bind to the rtTA allowing binding to the TRE and promoting transcription of the *Igα* and/or *Igβ* genes. Use of the tetracycline inducible system is exemplified in D.Y. Ho *et al.*, "Inducible gene expression from defective herpes simplex virus vectors using the

20 tetracycline-responsive promoter system," *Brain Res. Mol. Brain Res.* 41(1-2): 200-209, Sept. 5, 1996; Y. Yoshida *et al.*, "VSV-G-pseudotyped retroviral packaging through adenovirus-mediated inducible gene expression," *Biochem. Biophys. Res. Commun.* 232(2): 379-382, Mar. 17, 1997; A. Hoffman *et al.*, "Rapid retroviral delivery of tetracycline-inducible genes in a single autoregulatory cassette," *PNAS*, 93(11):

25 5185-5190, May, 28, 1996; and B. Massie *et al.*, "Inducible overexpression of a toxic protein by an adenovirus vector with a tetracycline-regulatable expression cassette," *J. Virol.* 72(3): 2289-2296, Mar. 1998, all of which are incorporated herein in their entireties by this reference.

Also provided by the present invention is a method of identifying a cell that produces a monoclonal antibody that recognizes a specific antigen comprising: a) immunizing a transgenic animal comprising B cells comprising a vector, wherein the vector comprises a nucleic acid encoding at least one surface-expressed antibody receptor selected from the group consisting of Ig α and Ig β ; b) isolating the B cells from the animal of step a); c) contacting the cells of step b) with the antigen, wherein the antigen binds to the monoclonal antibody to yield a detectable labeled cell; and d) isolating the detectably labeled cell, thus identifying a cell that produces a monoclonal antibody that recognizes a specific antigen.

The present invention also provides a hematopoietic stem cell comprising a vector, wherein the vector comprises a nucleic acid encoding at least one surface-expressed antibody receptor selected from the group consisting of Ig α and Ig β .

The present invention is more particularly described in the following examples which are intended as illustrative only since numerous modifications and variations therein will be apparent to those skilled in the art.

EXAMPLES

This invention shows that the lack of antibody receptors Ig α and/or Ig β , is the major limitation to surface presentation of antibody in hybridomas. The membrane form of antibody binds these two receptors through the membrane spanning domain that is on the C-terminus of the full-length heavy chain (mHC) as shown in **Figure 1**. Most myelomas have lost the ability to produce Ig α and/or Ig β , and the resulting hybridoma fusions no longer present surface mAb because they lack the the Ig α receptor or the Ig α and Ig β receptors (Kanavaros et al., 1995). Myelomas impart this lack of surface presentation of antibody to most hybridoma cell lines, even though many hybridomas are derived from early or mid-stage B-cells, which themselves present surface mAbs (Milcarek et al., 1996).

Engineering the constitutive expression of Igα and/or Igβ: The cDNAs encoding the two receptor sequences *Igα* and *Igβ* were PCR amplified from a mouse spleen cDNA library (Clontech). Restriction endonuclease cloning sites were added as part of the oligonucleotide primers used in the PCR amplification as shown in **Figure 2A** and the appropriate-sized PCR products were obtained (**Figure 2B**). The confirmed sequences of the PCR-amplified receptor for *Igα* and *Igβ* are shown in **Figures 3 and 4**, respectively. The PCR product containing *Igα* was digested with *HindIII* and *EcoRI* and cloned into the corresponding replacement region of the eukaryotic expression vector pcDNA3.1 (Neo) (Invitrogen, Inc.). The PCR product containing the *Igβ* sequence was digested with *HindIII* and *XhoI* and cloned into the corresponding replacement regions of the eukaryotic expression vector pcDNA3.1/Zeo (Invitrogen, Inc.). The structure of these two related pcDNA3.1 expression vectors are shown in **Figures 5 and 6**, respectively. The two vectors differ only in carrying resistance markers for Neomycin G418 and Zeocin, respectively. The resulting plasmids are termed p3.1Neo*Igα* and p3.1Zeo*Igβ*, respectively. Both pcDNA3.1 vectors express cloned sequences under the control of the strong constitutive CMV promoter and BGH terminator. Recombinant plasmid DNA was purified over an endotoxin free purification kit (Qiagen, Inc.) in preparation for transfection.

A well-characterized hybridoma cell line, HGS1 (fusion of Sp2/0 and a mouse B cell) was utilized (cloned line12G7). This line makes monoclonal antibodies to *E. coli* glutathione synthetase (GS) as described previously. The antibody reacts well with both a synthetic peptide designed from the GS sequence or the full-length GS protein.

The myeloma cell line Sp2/0 forms the standard fusion partner used for over a decade at University of Georgia's monoclonal facility in producing hybridomas. Sp2/0 and derived hybridomas are grown on RPMI medium (RPMI-1640, Sigma, Inc.) supplemented with 20% fetal Bovine serum (Atlanta Biologicals, Inc.) and grown at 37°C with 5% CO₂.

Optimizing transfection and selection: A constitutive β-galactosidase (β-gal) reporter plasmid (pcDNA3.1/lacZ, Invitrogen) was utilized to optimize and quantify

lipofection techniques on HGS1 hybridoma and Sp2/0 myeloma cell lines. Transfection was performed by mixing 6-8 μ l of LipofectAMINE reagent (Gibco BRL) with 1-2 μ g of plasmid DNA for 5 hr at 37°C in 1.0 ml of Opti-MEM I (GibcoBRL) reduced serum medium. Lipofection frequencies that occurred were relatively low, averaging approximately 30 transfectants per 500,000 cells, but were higher than previously reported for myeloma cells (Oi et al., 1983; Sun et al., 1991). The frequency of co-transfection of two DNAs was determined to average about 6-10 cells per 500,000. There was little difference between the frequency of transfecting or expressing linear or supercoiled plasmid DNA in several transfections, therefore, supercoiled DNA was used for subsequent experiments. Neomycin (Neo) (G418, Gibco BRL) and Zeocin (Invitrogen) kill curves were established on the same cells with 100% killing of control cells occurring over 7 days on 750 μ g/ml G418 and 750 mg/ml Zeocin. After this initial period of selection the G418 concentration remains the same, but the Zeocin concentration is reduced to 450 μ g/ml. Cells are grown under continuous selection.

Transfection and expression of Ig receptor genes: Receptor protein levels were assayed on Western blots of crude extracts resolved by SDS-PAGE. Rabbit polyclonal antibodies to the two receptors were provided by Dr. Linda Matsuuchi (Univ. Vancouver). Strong receptor expression is seen in the 30-40 kDa range for the spleen cell control (SC) as shown in **Figure 7**, while the higher molecular weight bands appear to be background. Using a double-drug selection for Neomycin and Zeocin isolated several independent and stably co-transfected cell lines (HGS1 α β 1-HGS1 α β 16) containing the two constructs p3.1NeoIg α and p3.1ZeolIg β were isolated. Several of these cell lines were examined for Ig α and Ig β expression on western blots. Two of five lines examined in one experiment, HGS1-Ig α β 2 and HGS1-Ig α β 5 (**Figure 7**) produced measurable levels of Ig α protein. This experiment also revealed that all cell lines examined produced significant amounts of Ig β with or without transfection with the pNeo3.1Ig α and p3.1ZeolIg β constructs. This background expression of Ig β was observed in myeloma line Sp2/0, the hybridoma line HGS1 (derived from a fusion

between Sp2/0 and a mouse B-cell), all lines derived from HGS1, and other hybridoma lines derived from Sp2/0.

Increased surface presentation of antibody in transfected hybridoma lines: The lines expressing high levels of Ig α from p3.1NeoIg α were examined for surface presentation of antibody in **Figure 8**. FITC-labeled sheep polyclonal anti-mouse antibody (Sigma) was used to measure the base level of mouse antibody on the surface of control cells, HGS1a. A low frequency of control cells present antibody with the typical result from several experiments being shown in **Figure 8A-B**. Remarkably, four of the six HGS1 α β cell lines transfected with both receptor plasmids (α β 6, α β 7, α β 9, α β 10) present large amounts of antibody on the surface of 100% of their cells as shown in **Figure 8C, D, F, and G**, respectively. A few cells in each field are out of focus, but a through focus examination of the field of cells reveals that 99% of the cells in each of the four populations present detectable levels of surface antibody. Clearly, examination of these cell populations reveals a significant increase in both the frequency of the cells that present antibody relative to the control cells and increases in the level of expression. Two of the G418 and Zeocin resistant transfected cell lines (α β 8, α β 11) showed no significant surface presentation of antibody (**Figure 8E and 8H**). Surprisingly, they presented less surface antibody, even than control HGS1 cells, with none of the 100-plus cells examined showing detectable surface expression.

Examining Ig receptor expression in hybridoma cells presenting surface antibody: The same cell samples examined in **Figure 8** were aliquoted and frozen for subsequent examination of receptor protein levels. Initial results comparing Ig α and Ig β receptor protein expression among the control HGS1 and transfected HGS1 α β cell lines are shown in **Figure 9 and 10**, respectively. Four of the Ig receptor plasmid transfected HGS1 α β cell lines express much more Ig α receptor compared to undetectable levels in control HGS1 cells. These cell lines expressing Ig α are the same lines showing 100% surface presentation of antibody in **Figure 8** (C α β 6, D α β 7, F α β 9, G α β 10). The α β 7 line showed significantly less Ig α protein expression than the three other lines showing strong surface expression. The α β 7 line showed surface expression of antibody in

essentially all cells examined (Figure 8D, $\alpha\beta 7$), but at lower intensity than the other three surface expressing lines, showing a direct quantitative relationship between Ig α levels and surface presentation. Two HGS1 $\alpha\beta$ cell lines showed no Ig α protein expression on the Western ($\alpha\beta 8$ and $\alpha\beta 11$), and these two showed even less surface presentation than the controls. It seems possible that some form of co-suppression of Ig α activity has occurred in these two negative lines. High levels of Ig β protein were detected all HGS1 derived cell lines examined and these levels did not correlate with surface presentation of antibody (Figure 10). Thus, increased surface antibody presentation on the HGS1 lines correlates directly and even semi-quantitatively with Ig α receptor protein expression.

Quantification of the mean fluorescence intensity of 50 cells in each population reveals that the transgenic Ig α expressing cells ($\alpha\beta 6$, $\alpha\beta 7$, $\alpha\beta 9$ and $\alpha\beta 10$) present about 5-times more antibody than control cells as shown in Figure 11A. A high percentage (60 to 80%) of strong Ig α expressing cells show a mean fluorescence intensity 3-times greater than the mean for non-Ig α expressing cells (Figure 11B). Only 0-6% of the control cells reach this level of intensity. The quantification of these data probably underestimated the actual increase in fluorescence of Ig α expressing cells. These assays are limited by the dynamic range of our instrumentation for measuring the most fluorescent transgenic cells (i.e., many Ig α cells are so bright they exceed the capacity of our instrumentation) and inability to assay all cells in a single focal plane. In addition, we see a low level of autofluorescence in labeled cells and weak background fluorescence in cells treated with an FITC labeled control antibody. This background may account for some of the fluorescence seen control cells (Figure 8A & B).

Additional experiments showed that Ig receptor overexpression and increased surface presentation of antibody did not prevent normal antibody secretion from hybridoma cells. Initial assays on cell supernatants from the eight lines examined in Figures 8-11 showed that each cell line still secreted significant level of mAbGS1 monoclonal antibody.

Myelomas The same genetic alteration utilized above on hybridoma cells is performed on a standard myeloma fusion partner Sp2/0. Sp2/0 is a myeloma cell line obtained from mice, *Mus musculus* (BALB/c). Sp2/0 is one of the founding myeloma cell lines used to make hybridoma fusions (Fraser and Venter, 1980; Greene et al., 1980; Hurwitz et al., 1980). First, Sp2/0 cells were co-transfected with the p3.1NeoIgx and p3.1ZeolIgβ constructs and selected for G418 and Zeocin resistance, to produce new cell lines Sp2αβ1, Sp2αβ2, etc. These Sp2αβ lines were characterized for Igα and Igβ receptor expression on western blots as shown in **Figure 12**. Lines Sp2αβ1 and Sp2αβ2, show strong Igα expression, and demonstrate that there is no post-transcriptional barrier to increasing receptor expression in myeloma cells. It appears that Igβ is already expressed at measurable levels in the control Sp2/0 control cells. These myeloma cell lines are ready to be fused with B cells in order to make hybridomas. Myeloma cells can be fused to a B cell or other antibody producing cell by methods standard in the art.

Fluorescent activated cell sorting further quantifies the increase in surface antibody presentation is linked to Igα expression: Total antibody on the surface of HGS1 cells and HGS1αβ10 cells was FITC labeled and the cells were sorted based on FITC fluorescence as shown in **Figure 13**. Comparison of panels A and B reveals that the mean fluorescence of the HGS1Igαβ10 cells is about 10-times greater than the fluorescence of HGS1 control cells (i.e., FL1 level of HGS1αβ10 cells in B is shifted about one log to the right of control HGS1 cells in A). Panel C examines the sorting of a mixture of these cells and shows a similar result. This difference in the mean level of fluorescence serves as an independent quantification of the effect of Igα expression to Figure 11. Increasing Igα expression results in an increase in surface presentation of mouse antibody in hybridoma cells. In sorting of normal hybridoma cell populations prepared freshly from splenic B-cells there would be none of the background fluorescence due to GS antibody presentation (right hand shoulder on the peak in A), because most cells would be making antibodies to antigens other than GS. In

conclusion, this clear increase in fluorescence due to Ig α expression demonstrates that DISH can be used in the direct selection of hybridomas.

Throughout this application, various publications are referenced. The disclosures of these publications in their entireties are hereby incorporated by reference into this application in order to more fully describe the state of the art to which this invention
 5 pertains.

REFERENCES

- 10 **Antczak, D.F.** (1982). Monoclonal antibodies: technology and potential use. *J Am Vet Med Assoc* **181**, 1005-10.
- Blattner, F.R., Plunkett, G., 3rd, Bloch, C.A., Perna, N.T., Burland, V., Riley, M., Collado-Vides, J., Glasner, J.D., Rode, C.K., Mayhew, G.F., Gregor, J., Davis, N.W., Kirkpatrick, H.A., Goeden, M.A., Rose, D.J., Mau, B., and Shao, Y.**
 15 (1997). The complete genome sequence of *Escherichia coli* K-12 [comment] [see comments]. *Science* **277**, 1453-74.
- Condon, C., Hourihane, S.L., Dang-Lawson, M., Escibano, J., and Matsuuchi, L.**
 (2000). Aberrant trafficking of the B cell receptor Ig-alpha beta subunit in a B lymphoma cell line. *J Immunol* **165**, 1427-37.
- 20 **Coursen, J.D., Bennett, W.P., Gollahon, L., Shay, J.W., and Harris, C.C.** (1997). Genomic instability and telomerase activity in human bronchial epithelial cells during immortalization by human papillomavirus-16 E6 and E7 genes. *Exp Cell Res* **235**, 245-53.
- Dickson, M.A., Hahn, W.C., Ino, Y., Ronfard, V., Wu, J.Y., Weinberg, R.A., Louis, D.N., Li, F.P., and Rheinwald, J.G.** (2000). Human keratinocytes that
 25 express hTERT and also bypass a p16(INK4a)-enforced mechanism that limits life span become immortal yet retain normal growth and differentiation characteristics. *Mol Cell Biol* **20**, 1436-47.

**Dunham, I., Shimizu, N., Roe, B.A., Chisoe, S., Hunt, A.R., Collins, J.E.,
Bruskiewich, R., Beare, D.M., Clamp, M., Smink, L.J., Ainscough, R.,
Almeida, J.P., Babbage, A., Bagguley, C., Bailey, J., Barlow, K., Bates, K.N.,
Beasley, O., Bird, C.P., Blakey, S., Bridgeman, A.M., Buck, D., Burgess, J.,
5 Burrill, W.D., O'Brien, K.P., and et al. (1999).** The DNA sequence of human
chromosome 22 [see comments] [published erratum appears in *Nature* 2000 Apr
20;404(6780):904]. *Nature* **402**, 489-95.

Edwards- Gilbert, G., and Milcarek, C. (1995). The binding of a subunit of the
general polyadenylation factor cleavage polyadenylation specificity factor
(CPSF) to polyadenylation sites changes during B cell development. *Nucleic Acids
Symposium Series* **33**, 229-233

Edwards-Gilbert, G., and Milcarek, C. (1995). Regulation of poly(A) site use during
mouse B-cell development involves a change in the binding of a general
polyadenylation factor in a B-cell stage-specific manner. *Molecular and Cellular
15 Biology* **15**, 6420-6429.

Edwards-Gilbert, G., Veraldi, K., and Milcarek, C. (1997). Alternative poly(A) site
selection in complex transcription units: means to an end? *Nucleic Acids Res* **25**,
2547-2561.

Flaspohler, J. A., Boczkowski, D., Hall, B. L., and Milcarek, C. (1995). The 3'-
20 untranslated region of membrane exon 2 from the gamma 2a immunoglobulin gene
contributes to efficient transcription termination. *Journal of Biological Chemistry*
270, 11903-11.

Flaspohler, J. A., and Milcarek., C. (1990). Myelomas and lymphomas expressing the
IgG2a H chain gene have similar transcription termination regions. *Journal of
25 Immunology* **144**, 2802-2810.

Fraser, C.M., and Venter, J.C. (1980). Monoclonal antibodies to beta-adrenergic
receptors: use in purification and molecular characterization of beta receptors. *Proc
Natl Acad Sci U S A* **77**, 7034-8.

- Galibert, F., Alexandraki, D., Baur, A., Boles, E., Chalwatzis, N., Chuat, J.C.,
Coster, F., Cziepluch, C., De Haan, M., Domdey, H., Durand, P., Entian, K.D.,
Gatius, M., Goffeau, A., Grivell, L.A., Hennemann, A., Herbert, C.J.,
Heumann, K., Hilger, F., Hollenberg, C.P., Huang, M.E., Jacq, C., Jauniaux,
5 J.C., Katsoulou, C., Karpfinger-Hartl, L., and et al. (1996). Complete nucleotide
sequence of *Saccharomyces cerevisiae* chromosome X. *Embo J* 15, 2031-49.
- Galli, G., Guise, J. W., McDevitt, M. A., Tucker, P. W., and Nevins, J. R. (1987).
Relative position and strengths of poly(A) sites as well as transcription termination
are critical to membrane vs secreted mu-chain expression during B-cell
10 development. *Genes & Dev* 1, 471-481.
- Genovese, C., Harrold, S., and Milcarek, C. (1991). Differential mRNA stabilities
affect mRNA levels in mutant mouse myeloma cells. *Somat Cell Mol Genet* 17, 69-
81.
- Genovese, C., and Milcarek, C. (1990). Increased half-life of mu immunoglobulin
15 mRNA during mouse B cell development increases its abundance. *Mol Immunol*
27, 733-43.
- Glennie, M.J., and Johnson, P.W. (2000). Clinical trials of antibody therapy.
Immunol Today 21, 403-10.
- Greenberg, R.A., O'Hagan, R.C., Deng, H., Xiao, Q., Hann, S.R., Adams, R.R.,
20 Lichtsteiner, S., Chin, L., Morin, G.B., and DePinho, R.A. (1999). Telomerase
reverse transcriptase gene is a direct target of c-Myc but is not functionally
equivalent in cellular transformation. *Oncogene* 18, 1219-26.
- Greene, G.L., Nolan, C., Engler, J.P., and Jensen, E.V. (1980). Monoclonal
antibodies to human estrogen receptor. *Proc Natl Acad Sci U S A* 77, 5115-9.
- 25 Guise, J., Lim, P., Yuan, D., and Tucker, P. (1988). Alternative expression of
secreted and membrane forms of immunoglobulin μ -chain is regulated by
transcriptional termination in stable plasmacytoma transfectants. *Journal of*
Immunology 140, 3988-3994.
- Hall, B. L., and Milcarek, C. (1989). Sequence and polyadenylation site determination

of murine immunoglobulin gamma 2a membrane 3' untranslated region. *Mol Immunol* 26, 819-826.

Hashimoto, S., Chiorazzi, N., and Gregersen, P.K. (1995). Alternative splicing of *CD79a (Ig-alpha/mb-1)* and *CD79b (Ig-beta/B29)* RNA transcripts in human B cells. *Mol Immunol* 32, 651-9.

Hattori, M., Fujiyama, A., Taylor, T.D., Watanabe, H., Yada, T., Park, H.S., Toyoda, A., Ishii, K., Totoki, Y., Choi, D.K., Soeda, E., Ohki, M., Takagi, T., Sakaki, Y., Taudien, S., Blechschmidt, K., Polley, A., Menzel, U., Delabar, J., Kumpf, K., Lehmann, R., Patterson, D., Reichwald, K., Rump, A., Schillhabel, M., and Schudy, A. (2000). The DNA sequence of human chromosome 21. The chromosome 21 mapping and sequencing consortium [see comments]. *Nature* 405, 311-9.

Hombach, J., Lottspeich, F., and Reth, M. (1990a). Identification of the genes encoding the IgM-alpha and Ig-beta components of the IgM antigen receptor complex by amino-terminal sequencing. *Eur J Immunol* 20, 2795-9.

Hombach, J., Tsubata, T., Leclercq, L., Stappert, H., and Reth, M. (1990b). Molecular components of the B-cell antigen receptor complex of the IgM class. *Nature* 343, 760-2.

Hurwitz, J.L., Coleclough, C., and Cebra, J.J. (1980). CH gene rearrangements in IgM-bearing B cells and in the normal splenic DNA component of hybridomas making different isotypes of antibody. *Cell* 22, 349-59.

Janeway, C. A., and Travers, P. (1994). *Immunobiology. The immune system in health and disease* (New York: Garland Publishing, Inc).

Kanavaros, P., Gaulard, P., Charlotte, F., Martin, N., Ducos, C., Lebezu, M., and Mason, D.Y. (1995). Discordant expression of immunoglobulin and its associated molecule mb-1/CD79a is frequently found in mediastinal large B cell lymphomas. *Am J Pathol* 146, 735-41.

Kandasamy, M.K., McKinney, E., and Meagher, R.B. (1999). The late pollen specific actins in angiosperms. *Plant J* 18, 681-691.

Kelly, D. E., and Perry, R. P. (1986). Transcriptional and post-transcriptional control of Ig mRNA production during B lymphocyte development. *Nucleic Acids Research* 14, 5431-5441.

Kim, H.S., Shin, J.Y., Yun, J.Y., Ahn, D.K., and Le, J.Y. (2001). Immortalization of human embryonic fibroblasts by overexpression of c- myc and simian virus 40 large T antigen. *Exp Mol Med* 33, 293-8.

Kiyono, T., Foster, S.A., Koop, J.I., McDougall, J.K., Galloway, D.A., and Klingelutz, A.J. (1998). Both Rb/p16INK4a inactivation and telomerase activity are required to immortalize human epithelial cells. *Nature* 396, 84-8.

Kohler, G., and Milstein, C. (1975). Continuous cultures of fused cells secreting antibody of predefined specificity. *Nature* 256, 495.

Kobrin, B. J., Milcarek, C., and Morrison, S. L. (1986). Sequences near the 3' secretion-specific polyadenylation site influence levels of secretion-specific and membrane-specific IgG2b mRNA in myeloma cells. *Molecular and Cellular Biology* 6, 1687-1697.

Konstantinos, N.S., and al., e. (1999). Use of monoclonal antibodies for the diagnosis and treatment of bladder cancer. *Hybridoma* 18, 219-224.

Lassman, C. R., and Milcarek, C. (1992). Regulated expression of the mouse γ 2b Ig H chain gene is influenced by polyA site order and strength. *J Immunol* 148, 2578-2585.

Lassman, C.R., Matis, S., Hall, B.L., Toppmeyer, D.L., and Milcarek, C. (1992). Plasma cell-regulated polyadenylation at the Ig gamma 2b secretion-specific poly(A) site. *J Immunol* 148, 1251-60.

Lebman, D. A., Park, M. J., Fatica, R., and Zhang, Z. (1992). Regulation of usage of membrane and secreted 3' termini of alpha mRNA differs from mu mRNA. *Journal of Immunology* 148, 3282-3289.

Li, Y., Kandasamy, M.K., and Meagher, R.B. (2001). Rapid isolation of monoclonal antibodies: monitoring enzymes in the phytochelatin synthesis pathway. *Plant Physiol* in press.

- Lockhart, D.J., and Winzeler, E.A.** (2000). Genomics, gene expression and DNA arrays. *Nature* **405**, 827-36.
- MacBeath, G., and Schreiber, S.L.** (2000). Printing proteins as microarrays for high-throughput function determination [see comments]. *Science* **289**, 1760-3
- 5 **Matis, S.A., Martincic, K., and Milcarek, C.** (1996). B-lineage regulated polyadenylation occurs on weak poly(A) sites regardless of sequence composition at the cleavage and downstream regions. *Nucleic Acids Res* **24**, 4684-92.
- McClelland, M., and Wilson, R.K.** (1998). Comparison of sample sequences of the *Salmonella typhi* genome to the sequence of the complete *Escherichia coli* K-12 genome. *Infect Immun* **66**, 4305-12.
- 10 **Meilhoc, E., Witttrup, K.D., and Bailey, J.E.** (1989). Application of flow cytometric measurement of surface IgG in kinetic analysis of monoclonal antibody synthesis and secretion by murine hybridoma cells. *J Immunol Methods* **121**, 167-74.
- Milcarek, C., and Hall, B.** (1985). Cell-specific expression of secreted versus
- 15 membrane forms of immunoglobulin gamma 2b mRNA involves selective use of alternate polyadenylation sites. *Mol Cell Biol* **5**, 2514-2520.
- Milcarek, C., Hartman, M., and Croll, S.** (1996). Changes in abundance of IgG 2a mRNA in the nucleus and cytoplasm of a murine B-lymphoma before and after fusion to a myeloma cell. *Mol Immunol* **33**, 691-701.
- 20 **Milcarek, C., Suda-Hartman, M., and Croll, S.C.** (1996). Changes in abundance of IgG 2a mRNA in the nucleus and cytoplasm of a murine B-lymphoma before and after fusion to a myeloma cell. *Mol Immunol* **33**, 691-701.
- Miller, R.A., Maloney, D.G., Warnke, R., and Levy, R.** (1982). Treatment of B-cell lymphoma with monoclonal anti-idiotypic antibody. *N Engl J Med* **306**, 517-22.
- 25 **Milstein, C.** (2000). With the benefit of hindsight. *Immunol Today* **21**, 359-64.
- Morio, T., Urushihara, H., Saito, T., Ugawa, Y., Mizuno, H., Yoshida, M., Yoshino, R., Mitra, B.N., Pi, M., Sato, T., Takemoto, K., Yasukawa, H., Williams, J., Maeda, M., Takeuchi, I., Ochiai, H., and Tanaka, Y.** (1998). The

Dictyostelium developmental cDNA project: generation and analysis of expressed sequence tags from the first-finger stage of development. *DNA Res* **5**, 335-40.

Mullner, S., Neumann, T., and Lottspeich, F. (1998). Proteomics--a new way for drug target discovery. *Arzneimittelforschung* **48**, 93-5.

5 **O'Reilly, L.A., Cullen, L., Moriishi, K., O'Connor, L., Huang, D.C., and Strasser, A.** (1998). Rapid hybridoma screening method for the identification of monoclonal antibodies to low-abundance cytoplasmic proteins. *Biotechniques* **25**, 824-30.

Oi, V.T., Morrison, S.L., Herzenberg, L.A., and Berg, P. (1983). Immunoglobulin gene expression in transformed lymphoid cells. *Proc Natl Acad Sci U S A* **80**, 825-9.

10 **Opitz, O.G., Suliman, Y., Hahn, W.C., Harada, H., Blum, H.E., and Rustgi, A.K.** (2001). Cyclin D1 overexpression and p53 inactivation immortalize primary oral keratinocytes by a telomerase-independent mechanism. *J Clin Invest* **108**, 725-32.

Pandey, A., and Mann, M. (2000). Proteomics to study genes and genomes. *Nature* 15 **405**, 837-46.

Pandey, A., Podtelejnikov, A.V., Blagoev, B., Bustelo, X.R., Mann, M., and Lodish, H.F. (2000). Analysis of receptor signaling pathways by mass spectrometry: identification of vav-2 as a substrate of the epidermal and platelet-derived growth factor receptors. *Proc Natl Acad Sci U S A* **97**, 179-84.

20 **Parks, D.R., Bryan, V.M., Oi, V.T., and Herzenberg, L.A.** (1979). Antigen-specific identification and cloning of hybridomas with a fluorescence-activated cell sorter. *Proc Natl Acad Sci U S A* **76**, 1962-6.

Persidis, A. (1998). Proteomics. *Nat Biotechnol* **16**, 393-4.

Price, C.P., and Newman, D.J. (1997). *Principles and practice of Immunoassay*,

25 **Richards, J.D., Gold, M.R., Hourihane, S.L., DeFranco, A.L., and Matsuuchi, L.** (1996). Reconstitution of B cell antigen receptor-induced signaling events in a nonlymphoid cell line by expressing the Syk protein-tyrosine kinase. *J Biol Chem* **271**, 6458-66.

Russo, I., Silver, A.R., Cuthbert, A.P., Griffin, D.K., Trott, D.A., and Newbold, R.F. (1998). A telomere-independent senescence mechanism is the sole barrier to Syrian hamster cell immortalization. *Oncogene* 17, 3417-26.

Sakaguchi, N., Kashiwamura, S., Kimoto, M., Thalmann, P., and Melchers, F. (1988). B lymphocyte lineage-restricted expression of mb-1, a gene with CD3-like structural properties. *Embo J* 7, 3457-64.

Sato, S., Nakamura, Y., Kaneko, T., Katoh, T., Asamizu, E., Kotani, H., and Tabata, S. (2000). Structural analysis of *Arabidopsis thaliana* chromosome 5. X. Sequence features of the regions of 3,076,755 bp covered by sixty P1 and TAC clones. *DNA Res* 7, 31-63.

Signals. (2000). Companies load up on magic bullets, *Signals Magazine* October, 1-9 (<http://www.recap.com/signalsma>).

Sun, L.K., Liou, R.S., Sun, N.C., Gossett, L.A., Sun, C., Davis, F.M., MacGlashan, D.W., Jr., and Chang, T.W. (1991). Transfectomas expressing both secreted and membrane-bound forms of chimeric IgE with anti-viral specificity. *J Immunol* 146, 199-205.

Yuan, D., and Tucker, P. W. (1984). Transcriptional regulation of the mu-delta heavy chain locus in normal murine B-lymphocytes. *J Exp Medicine* 160, 564-572.